

Circulating Allergen-reactive T Cells from Patients with Atopic Dermatitis and Allergic Contact Dermatitis Express the Skin-selective Homing Receptor, the Cutaneous Lymphocyte-associated Antigen

By Luis F. Santamaria Babi,* Louis J. Picker,†
M. Teresa Perez Soler,* Khristina Drzimalla,§ Peter Flohr,§
Kurt Blaser,* and Conrad Hauser*||

From the *Swiss Institute of Allergy and Asthma Research, CH-7270 Davos, Switzerland; the †Laboratory of Experimental Pathology, Department of Pathology, University of Texas Southwestern Medical School, Dallas, Texas 75235; §Klinik für Dermatologie und Allergie, CH-7270 Davos, Switzerland; and the ||Department of Dermatology, Hôpital Cantonal Universitaire, CH-1211 Geneva 14, Switzerland

Summary

The cutaneous lymphocyte-associated antigen (CLA) is the major T cell ligand for the vascular adhesion molecule E-selectin, and it has been proposed to be involved in the selective targeting of memory T cells reactive with skin-associated Ag to cutaneous inflammatory sites. To further investigate the relation of CLA and cutaneous T cell responses, we analyzed the CLA phenotype of circulating memory T cells in patients with allergic contact dermatitis and atopic dermatitis (AD) alone vs in patients manifesting bronchopulmonary atopy (asthma with or without AD) and nonallergic individuals. Significant T cell proliferative responses to Ni, a contact allergen, and to the house dust mite (HDM), an allergen to which sensitization is often observed in AD and/or asthma, was noted only in allergic and atopic individuals, respectively. When the minor circulating CLA⁺CD3⁺CD45RO⁺ subset was separated from the major CLA⁻CD3⁺CD45RO⁺ subpopulation in Ni-sensitive subjects, the Ni-dependent memory T cell response was largely confined to the CLA⁺ subset. A similar restriction of the T cell proliferative response to the CLA⁺ memory subset was observed for HDM in patients with AD alone. In HDM-sensitive patients with asthma with or without AD, however, the CLA⁻ subset exhibited a strong antigen-dependent proliferation, in contrast to patients with AD alone, whose CLA⁻ subset proliferated very weakly to HDM. In asthma with or without AD, the HDM-dependent proliferation slightly predominated in the CLA⁻ when compared to the CLA⁺ subset. The functional linkage between CLA expression and disease-associated T cell effector function in AD was also demonstrated by the finding that the circulating CLA⁺ T cell subset in AD patients, but not nonatopic controls, selectively showed both evidence of prior activation (human histocompatibility antigen-DR expression) and spontaneous production of interleukin 4 but not interferon- γ . Taken together, these observations demonstrate the correlation of CLA expression on circulating memory T cells and disease-associated memory T cell responses in cutaneous hypersensitivity, and they suggest the existence of mechanisms capable of sorting particular T cell Ag specificities and lymphokine patterns into homing receptor-defined memory subsets.

Recent investigation into the physiology of lymphocyte homing has revealed considerable heterogeneity in the ability of memory T cells to home to and localize within various lymphoid and extralymphoid tissues (1-4). In effect, the overall memory T cell population is divided into multiple subsets with varying, often tissue-specific, homing capabilities. One of the best-characterized "extralymphoid" T cell

homing specificities is the skin (5-7), which has long been postulated to represent a functionally distinct immune compartment and is a site of both localized T cell-mediated immune/inflammatory disease and localized T lineage malignancy (8, 9).

The interaction between circulating T cells and microvascular endothelial cells is a critical determinant of the homing

process, and we and others have proposed that tissue-selective T cell extravasation in the skin is mediated, at least in part, by the interaction between the cutaneous lymphocyte-associated antigen (CLA) on skin-homing T cells and its vascular counterreceptor, E-selectin, on inflamed superficial dermal postcapillary venules (10–17). In keeping with this hypothesis, CLA is expressed by a minor subset of CD45RO⁺ memory T cells in peripheral blood, by essentially all such cells in cutaneous sites of chronic inflammation, but by relatively few T cells in extracutaneous inflammatory sites (10, 14). Moreover, CLA is expressed by the malignant T cells of chronic-phase cutaneous T cell lymphoma (mycosis fungoides and Sézary syndrome), but not by non-skin-associated T cell lymphomas/leukemias (10, 15). CLA expression is regulated at the time of T cell activation by microenvironmental factors (including cytokines such as TGF- β 1 and IL-12) and has been demonstrated to be selectively induced on T cells undergoing the virgin to memory transition in skin-draining peripheral lymph nodes (PLN) (16 and Picker, L. J., unpublished data).

Since CLA is selectively associated with infiltrating T cells of the skin, it is possible that a similar association of CLA and skin-associated T cell functions exists in the blood. To investigate this possibility, we compared the distribution of disease-associated memory T cells in two skin-associated and MHC-restricted (18–20) immune responses to well-characterized Ag's—allergic contact dermatitis (CD) to Ni and atopic dermatitis (AD) associated with hypersensitivity to the house dust mite (HDM)—to that of T cells specific for tetanus toxoid (TT) in normal individuals, and to disease-associated T cells in patients who manifest bronchopulmonary atopy (asthma). Our results demonstrate the association of CLA expression on circulating memory T cells and cutaneous immune function in AD and CD, and they provide a model for the study of the mechanisms responsible for the selective distribution of the memory T cell repertoire among the different homing marker-defined T cell subsets.

Materials and Methods

Study Population. 14 patients with AD alone (mean age: 36 yr), 4 patients with AD plus asthma (mean age: 36 yr), 3 patients with asthma alone (mean age: 36 yr), 9 patients with Ni contact allergy (mean age: 35 yr), and 7 healthy controls (mean age: 36 yr) were analyzed in this study. All patients with AD fulfilled the criteria of Hanifin and Rajka (21). Atopics were skin prick test positive for HDM and presented HDM-specific IgE in serum. Ni-allergic CD patients showed positive epicutaneous patch test reactions to 0.5% NiSO₄ in Vaseline read at 48 and 72 h. Healthy controls had no history of atopy.

Antibodies and Ag's. mAb HECA-452 against CL available from L. J. Picker has previously been described (10). The mAbs anti-CD45RO (PE conjugated), anti-HLA-DR (PE conjugated), and anti-CD45RA were purchased from Immunotech (Marseille, France). The anti-CD16 mAb was obtained from the Central Laboratory of Blood Transfusion of the Red Cross (Amsterdam, The Netherlands) and anti-CD3-ECD from Coulter Corp. (Hialeah, FL). FITC-conjugated streptavidin was obtained from Tago, Inc. (Burlingame, CA). Goat anti-rat (H+L) biotin-SP-conjugated F(ab')₂ and goat anti-rat (H+L) biotin-SP-conjugated F(ab')₂ cou-

pled to FITC were purchased from Immunotech. Anti-CD14, anti-CD19, and streptavidin-conjugated magnetic microbeads were obtained from Miltenyi Biotech (Bergisch Gladbach, Germany).

HDM (*Dermatophagoides pteronyssinus*) extract was obtained from Haarlems Allergenen Laboratorium (Haarlem, The Netherlands). Purified *Dermatophagoides pteronyssinus* allergen (DerP1) was from Dr. H. Löwenstein (ALK, Horsholm, Denmark). Purified TT was obtained from the Swiss Serum and Vaccination Institute (Bern, Switzerland).

Preparation of CLA-enriched and CLA-depleted Memory T Cells. PBMC were isolated by Ficoll density gradient (Biochrom K.G., Berlin, Germany) centrifugation and resuspended in DMEM (Gibco, Basel, Switzerland) supplemented with 10 mM HEPES (Gibco), 5 mM EDTA (Fluka Chemie AG, Buchs, Switzerland), 2% FCS (Sera-Lab, Crawley Down, UK), 100 U/ml penicillin (Gibco), and 100 μ g/ml streptomycin (Gibco). Cells were mixed with anti-CD14 and anti-CD19 conjugated to magnetic microbeads (Miltenyi Biotech) and were passed through the magnetic activated cell separation system (MACS; Miltenyi Biotech) to obtain autologous APC. The CD14- and CD19-depleted cells were incubated with anti-CD45RA and anti-CD16 for 30 min on ice. Washed cells were mixed together with goat anti-mouse-conjugated magnetic microbeads to negatively select a population of CD45RO⁺ T cells. CD45RO⁺ T cells were incubated with anti-CLA HECA-452 mAb for 30 min on ice. Washed cells were then incubated with biotin-conjugated goat anti-rat (H+L) for 30 min on ice. Finally, washed cells were mixed with streptavidin conjugated to magnetic microbeads and were passed through the MACS. Both nonmagnetic (CLA⁻) and magnetic (CLA⁺) cells were recovered by sequential elution.

Proliferation Assays. The CLA-enriched and CLA-depleted memory T cells (5×10^4 cells per well) were incubated with 700 R irradiated autologous APC (10^4 cells per well) and with or without HDM (30 μ g/ml), DerP1 (10 μ g/ml), or NiSO₄ (20 μ g/ml) in 250 μ l of RPMI 1640 containing 10% complement-inactivated autologous serum, 2 mM L-glutamine (Gibco), 100 U/ml penicillin, and 100 μ g/ml streptomycin in 96-well U-bottom plates (Falcon Labware, Becton Dickinson and Co., Cockeysville, MD). The cultures were pulsed with 1 μ Ci of [³H]thymidine (DuPont/New England Nuclear, Boston, MA) after 5 d, and the incorporation of the labeled nucleotide was determined after 16 h.

Cell Staining and Flow Cytometry. Five-parameter analysis was performed on an Epics Profile (Coulter Corp.) installed with a power pack and equipped with a 488-nm argon laser. Fluorescence 1 was measured using a 525-nm band pass filter; fluorescence 2 was measured with a band pass filter of 575 nm, and fluorescence 3 was measured with a band pass filter of 635 nm. Typically, 10^5 cells were sequentially stained with HECA-452, goat anti-rat (H+L) conjugated to FITC, CD3-ECD, and CD45RO-PE, or HLA-DR-PE. Stained cells were fixed in 2% paraformaldehyde in PBS and stored at 4°C until they were analyzed. As a negative control, cells stained with FITC-conjugated goat anti-rat IgM were used. Cells were analyzed using the Gate Way software (Coulter Corp.).

IL4 and IFN- γ Assays. mAbs to IFN- γ and IL-4 were kindly provided by Dr. S. Alkan and Dr. C. Heusser, respectively (both from Ciba-Geigy, Basel, Switzerland), and ELISA assays were performed as described (22).

Statistical Analysis. Data were analyzed with the Mann-Whitney U test. The deviations shown in the results section represent 1 SEM.

Results

As delayed hypersensitivity to Ni manifesting as CD is exclusively expressed in the skin, we investigated the in vitro

proliferative response of CLA⁺ and CLA⁻ T cells to this allergen that is recognized in the context of MHC class II molecules (19, 23). Significant T cell proliferation to Ni was only observed in individuals with a positive patch test to Ni (data not shown). In these sensitive individuals (mean: 10.1% CLA⁺ in CD3⁺ PBMC; range: 7.0–18.3%), the proliferative response to the allergen was exclusively confined to the CLA⁺ circulating memory T cell subset in seven of nine subjects and nearly exclusively confined to this subset in the other two subjects (Fig. 1 A; $p = 0.0025$ for the comparison of CLA⁺ and ⁻ subsets for the overall group of nine subjects). In contrast, the proliferative response to TT was not confined to CLA⁺ memory T cells in five healthy subjects (mean: 12.0% CLA⁺ in CD3⁺ PBMC; range: 7.1–20.3%) and in four Ni-sensitive individuals (Fig. 1 B), demonstrating that not all antigen-induced memory T cell responses consistently predominate in the CLA⁺ subset.

AD is another skin-based T cell-mediated chronic hypersensitivity disorder in which responsible allergens can sometimes be identified. Unlike CD to Ni, however, atopic individuals may manifest hypersensitivity responses to a particular allergen in skin alone (AD), in an extracutaneous tissue alone

(primarily the lung, as in asthma), or in multiple tissues (either simultaneously in skin and lung or often at different times). The physiologic factors responsible for determining these tissue “preferences” of disease expression are unknown, but they may reflect differences in the regulation of homing receptor expression on developing memory T cells. To test this hypothesis, we identified a group of atopic individuals with demonstrable hypersensitivity and MHC-restricted (18, 20, 24–26) T cell reactivity to HDM who manifested atopic symptoms in the skin alone ($n = 7$) or in the lung alone or with the skin ($n = 7$). As shown in Fig. 2 A, memory T cell proliferative responses to HDM and DerP1 were preferentially detected in CLA⁺ T cells in six out of seven patients with AD alone ($p = 0.006$; mean: 7.3% CLA⁺ in CD3⁺ PBMC; range: 2.5–11.6%). In contrast, in those individuals manifesting asthma alone (mean: 8.2% CLA⁺ in CD3⁺ PBMC; range: 3.4–13.0%) or asthma with AD (mean: 14.3% CLA⁺ in CD3⁺ PBMC; range: 2.9–22.4%), the CLA⁻ subset exhibited a stronger HDM-dependent proliferative response than the CLA⁺ subset in patients with AD alone ($p = 0.0045$; Fig. 2). In all the patients with asthma




Figure 1. The Ni-induced proliferative response of circulating T cells from subjects with Ni sensitivity is largely confined to the CLA⁺ subset. CLA⁺ (solid bars) and CLA⁻ CD45RO⁺ (open bars) T cells from individuals with contact sensitivity to Ni (A) or sensitivity to TT (B) were incubated with or without Ni (A) or TT (B) in the presence of autologous CD14⁺CD19⁺ cells. Individuals tested with TT were either healthy controls or individuals with Ni sensitivity (S.U., S.C., H.H.). cpm >0 represent cpm in the presence of Ag minus cpm from cultures without Ag; cpm <0 represent cpm from cultures without Ag.




Figure 2. The HDM- and DerP1-induced proliferative response of circulating T cells from patients with AD is largely confined to the CLA⁺ subset. CLA⁺ (solid bars) and CLA⁻ CD45RO⁺ (open bars) T cells from patients with AD alone (A) and patients with asthma with (S.I., T.C., M.A., E.R.) or without AD (T.S., W.R., E.W.) (B) were incubated with or without HDM (DPT) or DerP1 in the presence of autologous CD14⁺CD19⁺ cells. The proliferative response was determined after 5 d. cpm >0 represent cpm in the presence of Ag minus cpm from cultures without Ag; cpm <0 represent cpm from cultures without Ag.

with or without AD, there was a slightly higher proliferative response to HDM in the CLA⁻ subset.

To further investigate the linkage between disease-associated T cells in AD and CLA expression, we asked whether patients with active AD may have increased numbers of circulating activated T cells, and if so, whether these cells expressed CLA. As shown in Fig. 3 A, patients with active AD did indeed demonstrate a significantly higher number of cells expressing the activation marker HLA-DR within the circulating CLA⁺ T cell subset as compared to the CLA⁻ subset (mean: 6.18 ± 1.6%, 1 SEM, vs. 1.56 ± 0.6, $p = 0.0035$). This is in contrast to the situation in normal

individuals, who demonstrate much lower overall HLA-DR expression and no preferential expression of the CLA⁺ T cell subset (Fig. 3 B). Moreover, in AD subjects (Fig. 3 C), unlike in normal control subjects (data not shown), a high level of spontaneous IL-4 production was observed after *in vitro* incubation of the CLA⁺, but not CLA⁻, memory T cell subset. Furthermore, IFN- γ was not detected in these subjects in either T cell subset (data not shown). Given that IL-4-producing (Th2-type) T cells have been strongly implicated in the pathogenesis of AD (27–31), these results suggest that when dermatitis is the primary presentation of atopy, disease-associated effector T cells are primarily in the CLA⁺ memory subset.

Discussion

AD and allergic CD are two independent forms of cutaneous allergy that are associated with an allergen-dependent T cell response (32). In animal models, it has clearly been documented that allergic CD is T cell dependent and that the allergen-specific T cells migrate into the site of allergen application (33). There they may induce the clinical features of dermatitis, which include swelling, erythema, and secondary infiltration, first by granulocytes and later by mononuclear cells. There is evidence that allergic CD is a type 1 Th response (34). Although it has not formally been shown that type 2 Th are responsible for AD, type 2 Th clones with specificity for atopy-related Ag's can be derived from the skin of patients with AD (27–29). In addition, so-called late-phase cutaneous reactions in atopics contain many T cells and exhibit positive signals for type 2 Th lymphokines by *in situ* hybridization (30). Although there is presently no animal model for AD, activated type 2 Th injected into the skin can induce inflammation that is dependent on IL-4 (31). Similarly, type 2 Th cytokines are found in bronchoalveolar cells from patients with allergic asthma (35, 36). Such IL-4- and IL-5-producing T cells in the bronchial wall may be responsible for the recruitment of eosinophils and other proinflammatory cells, as well as for the late inflammatory phases of asthma.

In addition to the skin-selective expression of CLA in organ-infiltrating T cells, evidence linking CLA expression of memory T cells to skin-selective homing includes (a) the pattern of expression of other tissue-selective homing receptors on the CLA⁺ T subset (high L-selectin; low $\alpha_4\beta_7$ - and $\alpha_e\beta_7$ -integrins; 14); (b) the high expression of CLA on allergen-specific T cell clones from an AD patient vs the lack of CLA on non-Ag-specific blood-derived clones from the same patient (13); and (c) the selective up-regulation of CLA on T cells undergoing the virgin to memory transition in skin-draining PLNs vs mucosal secondary lymphoid tissues (16). Here we provide further evidence for the association of CLA and skin immune responses by demonstrating that the circulating Ag-specific memory T cells in subjects with skin-restricted chronic inflammation conditions are highly restricted to the CLA⁺ subset. Moreover, in AD patients, circulating HLA-DR-expressing and “spontaneously” IL-4-producing T cells were selectively enriched in the CLA⁺




Figure 3. CLA⁺ circulating T cells from patients with AD contain an increased percentage of HLA-DR⁺ cells and spontaneously release increased amounts of IL-4. PBMCs from patients with AD (A) and healthy controls (B) were stained with mAb to CD3, HLA-DR, and CLA. CD3⁺ cells were gated and the percentage of HLA-DR⁺ cells within the CLA⁺ (solid bars) and CLA⁻ (open bars) subsets was determined. Memory T cells were isolated from PBMCs of patients with AD and cultured with CD14⁺ and CD19⁺ cells. IL-4 was determined in the 48-h supernatant (C).

subset. In contrast, the CLA⁻ subset was the major source of HDM reactivity in HDM-sensitive asthmatic individuals. Taken together, our observations are compatible with the skin-homing function of CLA, and they suggest the existence of mechanisms that are capable of selectively targeting memory T cells with a particular Ag reactivity and lymphokine pattern into a distinct homing marker-defined subset.

We have previously proposed that homing receptor expression, like effector functions such as cytokine synthesis potential (37), is regulated at the time of T cell activation, initially during the virgin to memory transition in secondary lymphoid tissue, and perhaps also at subsequent "reactivations" in secondary lymphoid tissues or extralymphoid effector sites (16). However, similar to what has been described for cytokine production, the specific regulatory elements directing homing function *in vivo* are likely to be complex and thus may not always correlate with anatomic site of antigen deposition alone. Nevertheless, this hypothesis would predict that T cells reactive with an Ag that consistently is encountered by the immune system in CLA-inducing microenvironment would, over

time, preferentially accumulate in the skin-homing CLA^{high}, L-selectin⁺, $\alpha_4\beta_7$ -integrin⁻ memory T cell subset.

Contact allergens such as Ni would clearly fit into the category of tissue (e.g., skin)-associated Ag's, and our finding that Ni-reactive memory T cells are almost exclusively in the CLA⁺ subset supports this hypothesis. It is much less clear how Ag's such as HDM interact with the immune system, and indeed the varied organs of atopy manifestations in different patients and in the same patient at different times suggest variability of the microenvironment in which specific T cells of atopics encounter HDM Ag. Clearly, however, in some individuals with atopic manifestations restricted to the skin, HDM-specific T cells are preferentially in the CLA⁺ memory subset. It remains to be determined in these patients whether this preferential CLA induction on HDM-reactive memory T cells results from HDM preferentially entering the body via cutaneous routes or from genetic or other regulatory influences promoting CLA induction in non-skin-associated microenvironments.

This work was supported by the Swiss National Foundation for Scientific Research grants 31-29967.90 and 31-39430.93, and by the EMDO Stiftung. Dr. L. F. Santamaria Babi's contribution was supported by the Walter Siegenthaler Stipendium. Dr. Picker's contribution was supported by National Institutes of Health grant AI-31545.

Address correspondence to Dr. Luis F. Santamaria Babi, Swiss Institute for Allergy and Asthma Research, Obere Strasse 22, 7270 Davos, Switzerland.

Received for publication 16 November 1994 and in revised form 22 December 1994.

References

1. Picker, L.J., and E.C. Butcher. 1992. Physiological and molecular mechanisms of lymphocyte homing. *Annu. Rev. Immunol.* 10:561-591.
2. Springer, T.A. 1994. Traffic signals for lymphocyte recirculation and leukocyte emigration: the multistep paradigm. *Cell.* 76:301-314.
3. Picker, L.J. 1994. Control of lymphocyte homing. *Curr. Opin. Immunol.* 6:394-406.
4. Mackay, C.R. 1993. Homing of naive, memory and effector lymphocytes. *Curr. Opin. Immunol.* 5:423-427.
5. Mackay, C.R., W. Marston, and L. Dudler. 1992. Altered patterns of T cell migration through lymph nodes and skin following antigen challenge. *Eur. J. Immunol.* 22:2205-2210.
6. Rose, M.L., D.M. Parrott, and R.G. Bruce. 1976. Migration of lymphoblasts to the small intestine. II. Divergent migration of mesenteric and peripheral immunoblasts to sites of inflammation in the mouse. *Cell. Immunol.* 27:36-46.
7. Issekutz, T.B., W. Chin, and J.B. Hay. 1980. Lymphocyte traffic through granulomas: differences in the recovery of indium-111-labeled lymphocytes in afferent and efferent lymph. *Cell. Immunol.* 54:79-86.
8. Bos, J.D., I. Zonneveld, P.K. Das, S.R. Krieg, C.M. van der Loos, and M.L. Kapsenberg. 1987. The skin immune system (SIS): distribution and immunophenotype of lymphocyte subpopulations in normal human skin. *J. Invest. Dermatol.* 88:569-573.
9. Edelson, R.L. 1980. Cutaneous T cell lymphoma: mycosis fungoides, Sézary syndrome, and other variants. *Am. Acad. Dermatol.* 2:89-106.
10. Picker, L.J., S.A. Michie, L.S. Rott, and E.C. Butcher. 1990. A unique phenotype of skin-associated lymphocytes in humans: preferential expression of the HECA-452 epitope by benign and malignant T-cells at cutaneous sites. *Am. J. Pathol.* 136:1053-1068.
11. Picker, L.J., T.K. Kishimoto, C.W. Smith, R.A. Warnock, and E.C. Butcher. 1991. ELAM-1 is an adhesion molecule for skin-homing T cells. *Nature (Lond.)* 349:796-799.
12. Berg, E.L., T. Yoshino, L.S. Rott, M.K. Robinson, R.A. Warnock, T.K. Kishimoto, L.J. Picker, and E.C. Butcher. 1991. The cutaneous lymphocyte antigen is a skin lymphocyte homing receptor for the vascular lectin endothelial cell-leukocyte adhesion molecule 1. *J. Exp. Med.* 174:1461-1466.
13. Rossiter, H., G.C. Mudde, F. van Reijssen, F. Kalthoff, C.A.F.M. Bruijnzeel-Koomen, L.J. Picker, and T.S. Kupper. 1994. Disease-related T cells from atopic skin express cutaneous lymphocyte antigen and sialyl Lewis X determinants, and bind to both E-selectin and P-selectin. *Eur. J. Immunol.* 24:205-210.
14. Picker, L.J., R.J. Martin, A. Trumble, L.S. Newman, P.A. Collins, P.R. Bergstresser, and D.Y.M. Leung. 1994. Differential expression of lymphocyte homing receptors by human

- memory/effector T cells in pulmonary vs. cutaneous immune effector sites. *Eur. J. Immunol.* 24:1269–1277.
15. Heald, P.W., S.-L. Yan, R.L. Edelson, R. Tigelaar, and L.J. Picker. 1993. Skin-selective lymphocyte homing mechanisms in the pathogenesis of leukemic cutaneous T-cell lymphoma. *J. Invest. Dermatol.* 101:222–226.
 16. Picker, L.J., J.R. Treer, B. Ferguson-Darnell, P.A. Collins, P.R. Bergstresser, and L.W.M.M. Terstappen. 1993. Control of lymphocyte recirculation in man. II. Differential regulation of the cutaneous lymphocyte-associated antigen, a tissue-selective homing receptor for skin-homing T cells. *J. Immunol.* 150:1122–1136.
 17. Santamaria, L.F., R. Moser, M.T. Perez Soler, L.J. Picker, K. Blaser, and C. Hauser. 1995. The migration of skin-homing T cells across cytokine-activated human endothelial cell layers involves interaction of the cutaneous lymphocyte-associated antigen (CLA), the very late antigen-4 (VLA-4) and the lymphocyte function-associated antigen-1 (LFA-1). *J. Immunol.* 154: 1543–1550.
 18. Yssel, H., K.E. Johnson, P.V. Schneider, J. Wideman, A. Terr, R. Kastelein, and J.E. De Vries. 1992. T cell activation-inducing epitopes of the house dust mite allergen Der p1. Proliferation and lymphokine production patterns by Der p1-specific CD4⁺ T cell clones. *J. Immunol.* 148:738–745.
 19. Kapsenberg, M.L., T. Van der Pouw-Kraan, F.E. Stiekema, A. Schootemeijer, and J.D. Bos. 1988. Direct and indirect nickel-specific stimulation of T lymphocytes from patients with allergic contact dermatitis to nickel. *Eur. J. Immunol.* 18:977–982.
 20. Lin, K.L., K.H. Hsieh, J.K. Huang, and S.Y. Wang. 1992. Major histocompatibility complex (MHC) class II restriction, lymphokine production, and IgE regulation of house dust mite-specific T cell clones. *J. Clin. Immunol.* 12:271–280.
 21. Hanifin, J.M., and G. Rajka. 1980. Diagnostic features of atopic dermatitis. *Acta Derm. Venereol. (Stockh.)* 92:44–47.
 22. Brinkman, V., T. Geiger, S. Alkan, and C.H. Heusser. 1993. Interferon α increases the frequency of Interferon γ -producing human CD4⁺ T cells. *J. Exp. Med.* 178:1655–1663.
 23. Sinigaglia, F., D. Scheidegger, G. Garotta, P. Scheper, M. Pletscher, and A. Lanzavecchia. 1985. Isolation and characterization of Ni-specific T cell clones from patients with Ni-contact dermatitis. *J. Immunol.* 135:3929–3932.
 24. O'Hehir, R.E.O., and J.R. Lamb. 1991. The cellular and molecular basis of the human T cell response to *Dermatophagoides spp.* (house dust mite). *Parasite Immunol. (Oxf.)* 13:109–119.
 25. Lamb, J.R., and R.E. O'Hehir. 1989. HLA class II restriction specificity of *Dermatophagoides spp.* reactive T lymphocyte clones that support IgE synthesis. *Clin. Exp. Allergy* 19:389–393.
 26. O'Hehir, R.E., B. Mach, C. Berte, R. Greenlaw, J.-M. Tiercy, V. Bal, R.I. Lechler, J. Trowsdale, and J.R. Lamb. 1990. Direct evidence for a functional role of HLA-DRB1 and -DRB3 gene products in the recognition of *Dermatophagoides spp.* (house dust mite) by helper T lymphocytes. *Int. Immunol.* 2:885–892.
 27. Van der Heijden, F.L., E.A. Wierenga, J.D. Bos, and M.L. Kapsenberg. 1991. High frequency of IL-4-producing CD4⁺ allergen-specific T lymphocytes in atopic dermatitis lesional skin. *J. Invest. Dermatol.* 97:389–394.
 28. Sager, N., A. Feldmann, G. Schilling, P. Kreitsch, and C. Neumann. 1992. House dust-mite specific T cells in the skin of subjects with atopic dermatitis: frequency and lymphokine profile in the allergen patch test. *J. Allergy Clin. Immunol.* 89:801–810.
 29. Van Reijssen, F.C., C.A.F.M. Bruijnzeel-Koomen, F.S. Kalthoff, E. Maggi, S. Romagnani, J.K.T. Westland, and G.C. Mudde. 1992. Skin-derived aero-allergen specific T cell clones of Th2 phenotype in patients with atopic dermatitis. *J. Allergy Clin. Immunol.* 90:184–193.
 30. Kay, A.B., S. Ying, V. Varney, M. Gaga, S.R. Durham, R. Moqbel, A.J. Wardlaw, and Q. Hamid. 1991. Messenger RNA expression of the cytokine gene cluster interleukin 3 (IL-3), IL-4, IL-5, and granulocyte/macrophage colony-stimulating factor, in allergen-induced late-phase cutaneous reactions in atopic subjects. *J. Exp. Med.* 173:775–778.
 31. Müller, K.M., F. Jaunin, I. Masouyé, J.-H. Saurat, and C. Hauser. 1993. Th2 cells mediate IL-4-dependent local tissue inflammation. *J. Immunol.* 150:5576–5584.
 32. Kägi, M.K., B. Wüthrich, E. Montano, J. Barandun, K. Blaser, and C. Walker. 1994. Differential cytokine profiles in peripheral blood lymphocyte supernatants and skin biopsies from patients with different forms of atopic dermatitis, psoriasis and normal individuals. *Int. Arch. Allergy Immunol.* 103:332–340.
 33. Ferguson, T.A., and T.S. Kupper. 1993. Antigen-independent processes in antigen-specific immunity. A role for α , integrin. *J. Immunol.* 150:1172–1182.
 34. Kapsenberg, M.L., E.A. Wierenga, F.E.M. Stiekema, A.M.B.C. Tiggeleman, and J.D. Bos. 1992. Th1 lymphokine production profiles of Ni-specific CD4⁺ T-lymphocyte clones from Ni contact allergic and non-allergic individuals. *J. Invest. Dermatol.* 98:59–63.
 35. Walker, C., E. Bode, L. Boer, T.T. Hansel, K. Blaser, and J.-C. Virchow, Jr. 1992. Allergic and nonallergic asthmatics have distinct patterns of T-cell activation and cytokine production in peripheral blood and bronchoalveolar lavage. *Am. Rev. Respir. Dis.* 146:109–115.
 36. Walker, C., J.-C. Virchow, Jr., P.L.B. Bruijnzeel, and K. Blaser. 1991. T cell subsets and their soluble products regulate eosinophilia in allergic and nonallergic asthma. *J. Immunol.* 146:1829–1835.
 37. Paul, W.E., and R.A. Seder. 1994. Lymphocyte responses and cytokines. *Cell.* 76:241–251.