
Brief Definitive Report

**GRAFT-VERSUS-HOST-RELATED IMMUNOSUPPRESSION IS
INDUCED IN MIXED CHIMERAS BY ALLORESPONSES
AGAINST EITHER HOST OR DONOR
LYMPHOHEMATOPOIETIC CELLS**

BY MEGAN SYKES, MICHAEL A. SHEARD, AND DAVID H. SACHS

*From the Transplantation Biology Section, Immunology Branch, National Cancer Institute,
National Institutes of Health, Bethesda, Maryland 20892*

Graft-versus-host disease (GVHD) in man and animals is associated with immune dysfunction, which results in significant morbidity and mortality (1). However, the precise role of graft-versus-host (GVH) reactivity in producing immune dysfunction is difficult to evaluate, because ablation of host immunity is necessary before GVHD can be induced. Examination of the effects of GVH reactions without reciprocal reactions of host against donor have involved the use of recipients that are not immunocompetent, such as neonatal mice (2), and adult animals in which the immune system has been artificially ablated by lethal or sublethal irradiation (3-6). In addition, GVH reactions have been induced in recipients that are initially immunocompetent but specifically tolerant of donor inocula. For example, GVHD has been induced by injection of donor lymphocytes into adult rodents neonatally tolerized to donor antigen (7, 8), and the effects of GVH inocula on immune function have been extensively studied in immunocompetent F₁ animals inoculated with parental lymphocytes (9-13). Loss of immune responsiveness after inoculation of F₁ recipients with parental lymphocytes has been variably attributed to suppressor cells (9-11), destruction of recipient lymphocytes (11, 14, 15), and thymic injury (9, 12, 13). However, the contribution of each of these mechanisms has not been well defined. We have now studied the effects of GVH reactions on immune responsiveness in animals with otherwise intact immune systems, in which tolerance of recipients to the donor inoculum is not due to a genetic mechanism, but has instead been induced by lethal irradiation followed by reconstitution with T cell-depleted (TCD) bone marrow (BM). Comparison of the effects of GVH inocula in fully allogeneic vs. mixed allogeneic chimeras as well as examination of the effects of host vs. graft (HVG) inocula have allowed examination of the role of recipient lymphohematopoietic cells, in distinction from the remainder of the animal's tissues, in inducing the immune abnormalities associated with GVH reactions in vivo.

Materials and Methods

Radiation bone marrow chimeras were prepared as previously described (16, 17). Briefly, recipient C57BL/10SnJ (B10) or B10.D2nSn (B10.D2) mice, 12-14 wk old, received lethal whole body irradiation and were reconstituted with allogeneic (B10 or B10.D2) with or without syngeneic (recipient strain) TCD bone marrow. Appropriate levels of chimerism were ascer-

tained in all chimeric responders by H-2 phenotyping of PBL by fluorescence microfluorimetry performed 6–10 wk after bone marrow transplantation (BMT) (18). Animals receiving 5×10^6 TCD syngeneic plus 1.5×10^7 TCD allogeneic BM cells reconstituted as stable mixed chimeras, demonstrating 25–75% allogeneic lymphoid chimerism, and are referred to here as “mixed” chimeras. Animals reconstituted with 1.5×10^7 TCD allogeneic cells alone are designated “allogeneic” chimeras and demonstrated 94–99% allogeneic chimerism. Previous studies have established the presence of specific tolerance to donor and host antigen and permanent, stable chimerism in such animals (16, 17). GVH inocula consisting of $2\text{--}3 \times 10^7$ splenocytes from normal donor-strain mice were administered via the lateral tail vein 10–22 wk after BMT. These recipients, as well as uninjected control chimeras, were killed 12 d after inoculation, and cell-mediated lympholysis (CML) responses of their splenocytes were assessed as previously described (19).

Results and Discussion

GVH inocula consisting of $2\text{--}3 \times 10^7$ splenocytes from normal donor-strain mice were administered to long-term stable chimeras. Fig. 1 (*upper panels*) shows typical CML responses of a long-term B10 + B10.D2 \rightarrow B10 mixed allogeneic chimera 12 d after inoculation with B10.D2 splenocytes. The uninjected animal was unresponsive to donor (B10.D2) and host (B10) antigen, but responded normally to third-




FIGURE 1. Cell-mediated lympholysis (CML) responses of chimeras against (A) host (B10), (B) donor (B10.D2), and (C) third-party (B10.BR) alloantigen 12 d after inoculation with normal donor-strain (B10.D2) splenocytes (□) or in unmanipulated control chimeras (○). Positive control responses (■) of normal (A) B10.D2 or (B) B10 splenocytes as well as negative control responses (●) of normal (A) B10, (B) B10.D2, or (C) B10.BR splenocytes are also shown. Irradiated stimulator cells and target blasts (stimulated with Con A for 2 d in culture) were obtained from normal B10 mice in A, from B10.D2 mice in B, and from B10.BR mice in C. Upper panels show responses of mixed chimeras (B10 + B10.D2 \rightarrow B10), and lower panels show responses of an allogeneic chimera (B10.D2 \rightarrow B10) given an identical B10.D2 splenocyte inoculum at the same time, as well as those of an unmanipulated allogeneic chimera.

party (B10.BR) alloantigen. The injected chimera, in contrast, was severely hyporesponsive towards third-party alloantigen. Similar anti-third-party CML hyporesponsiveness has been observed in 17 of 17 mixed chimeras tested from 10 to 28 d after administration of GVH inocula. Anti-third-party alloresponsiveness was lost between 5 and 10 d after inoculation, and recovered in most animals by 7–12 wk (data not shown). Similar to uninjected control chimeras, the injected mixed chimera shown in Fig. 1 was unresponsive to donor antigen. It did, however, demonstrate slight CML reactivity towards host antigen, which was a common but variable finding among injected mixed chimeras.

Alloresponses were studied in B10.D2 \rightarrow B10 (allogeneic) chimeras after administration of similar GVH inocula, and compared with those of allogeneic chimeras not receiving GVH inocula. Unlike mixed chimeras given identical inocula in the same experiments, allogeneic chimeras retained CML responsiveness to third-party alloantigen (Fig. 1 C, *lower panel*). Such responsiveness was maintained at all time points tested (5, 10, 12, 13, 21, and 28 d after GVH inoculum). Injected allogeneic chimeras also remained unresponsive to donor antigen, and some demonstrated slight CML reactivity towards host antigen.

Therefore, these studies demonstrated that mixed chimeras, when inoculated with nontolerant donor-strain cells, lost the ability to generate CTL in response to third-party alloantigen. These results are similar to those described in F₁ recipients, in which administration of parental cells recognizing F₁ class I plus class II alloantigens results in loss of CML responses to third-party alloantigen (10, 11, 14, 20). In contrast, when donor-strain lymphocytes were injected into allogeneic chimeras, CML responses of recipient chimeras to third-party alloantigen were not significantly affected. These findings suggested that the development of GVH-related immunodeficiency might depend on the presence of substantial numbers of lymphohematopoietic cells expressing the recipient phenotype, and that the recognition of such antigen on other somatic cells of the host was not sufficient to induce immunodeficiency. We reasoned that if this were true, then the presence of a one-way response to donor-type lymphohematopoietic cells might also be sufficient to lead to immune hyporesponsiveness, in this case induced by administration of nontolerant recipient-strain lymphocytes to mixed chimeras (i.e., HVG inocula). Such mixed chimeras would provide donor BM-derived lymphohematopoietic cells as a source of stimulating antigen, and should be tolerant of recipient antigen (16, 17), and therefore, should not react against injected recipient-strain lymphocytes.

To test this hypothesis, 3×10^7 recipient-strain splenocytes were injected into long-term B10 + B10.D2 \rightarrow B10 and B10 + B10.D2 \rightarrow B10.D2 mixed chimeras. As a control, additional mixed chimeras of both types received GVH inocula (3×10^7 donor-strain, B10 or B10.D2 splenocytes) in the same experiment. Representative anti-third-party alloresponses from such animals 12 d after inoculation are shown in Fig. 2. Injection of either B10 or B10.D2 splenocytes into mixed chimeras of either B10 or B10.D2 recipient strains resulted in marked hyporesponsiveness to third-party alloantigen, irrespective of whether the splenocytes represented a GVH or an HVG inoculum. Uninjected control animals, in contrast, remained responsive to third-party alloantigen. Thus, the presence of a one-way alloresponse to either donor or host antigen on lymphohematopoietic cells in vivo was sufficient to induce hyporesponsiveness to third-party alloantigen, indicating that induction of unresponsiveness does not specifically require a GVH reaction. It therefore seems likely that induction of GVH-related immunodeficiency described in parent into F₁ systems (9–12, 15, 20)




FIGURE 2. Effects of GVH and HVG inocula on anti-third-party CML responses in mixed allogeneic chimeras. Splenocytes were removed from mixed B10 + B10.D2 \rightarrow B10 (left panel) and B10 + B10.D2 \rightarrow B10.D2 (right panel) chimeras 13 d after inoculation with B10 (\blacktriangle) or B10.D2 (\circ) lymphocytes, or after no injection (\bullet). Curves represent percent specific lysis at various responder/target ratios of chimeric splenocytes on B10.BR Con A blasts after incubation for 5 d with irradiated B10.BR stimulator cells (see legend to Fig. 1). Similar results were obtained in three independent experiments.

might not require a generalized GVH reaction, but instead might reflect a one-way immune response in vivo to host lymphohematopoietic components.

These results are incompatible with one possible explanation for the immune dysfunction observed in F_1 animals inoculated with parental lymphocytes, namely the destruction of recipient T cells by the donor inoculum (11, 14) with selective overgrowth of host-reactive donor-derived T cells (15). Mixed chimeras contain T cells derived from both donor and recipient strain BM, and preliminary data indicate that both types of T cells are capable of mediating CTL responses (unpublished data). Therefore, those donor- or recipient- BM-derived T cells that are capable of mediating immune responses against third-party alloantigen but that are not potential targets of GVH- or HVG-reactive cells, respectively, should persist after GVH or HVG inoculation. The apparent inability of these cells to respond to third-party alloantigen suggests that they may in some way be inactivated as a result of the GVH reaction. Preliminary studies suggest this may be due to the induction of a G10-adherent suppressor cell population (Sykes et al., unpublished data).

Complete repopulation with allogeneic cells occurred in most mixed chimeras by 30–60 d after administration of GVH inocula. Although such repopulation is evidence for the occurrence of antihost aggressive activity in vivo, these animals survived long-term with no clinical evidence of GVHD. The numbers of donor-strain lymphocytes administered in GVH inocula in these experiments greatly exceeded those required to produce lethal GVHD in fresh irradiated animals (21). It is therefore possible that suppressive activity producing the immune hyporesponsiveness described here may also protect recipients from GVHD. However, tolerance to donor alloantigen is rapidly broken in recipients of similar HVG inocula to those used here (22), indicating that the immunosuppression resulting from such inocula is not effective in preventing the alloreactivity associated with graft rejection. Alternative explanations for the absence of GVHD in mixed chimeric recipients of GVH inocula, therefore, also deserve consideration. Induction of clinical GVHD may require factors in addition to one-way immune responses against host lymphohematopoietic cells in vivo, such as the acute tissue injury and infections that may follow lethal irradiation (9). It is also possible that a form of alloresistance against donor lymphoid cells exists in mixed chimeras similar to that which protects F_1 mice from the induction of GVH-related effects by parental lymphocytes of certain strains (20).

In summary, the studies presented here suggest a requirement for lymphohematopoietic cells bearing host antigen in the induction of GVH-related immunosup-

pression in radiation chimeras. A similar defect in immune responses was found in mixed chimeras given inocula that attack donor and not recipient antigens. These findings are most consistent with the interpretation that the immunodeficiency associated with GVH reactions is due to the development of suppressive activity as a consequence of a strong one-way alloresponse directed against lymphohematopoietic cells in vivo, whether this be an HVG or a GVH response. The presence of such immune suppression in mixed chimeras did not lead to death from infection, and did not appear to prevent the rapid loss of tolerance to donor antigen in animals given HVG inocula (22). The apparent discrepancy between the tolerance-breaking studies (22) and those reported here might either be due to differences in the time courses of the in vitro and in vivo phenomena, or might indicate that in vitro unresponsiveness does not accurately reflect the in vivo immune status of such recipients. Further studies will be required to elucidate the in vivo consequences, if any, of the immune unresponsiveness observed in vitro after administration of HVG and GVH inocula. One potential consequence to be addressed is the possibility that in vivo reactivity against injected lymphohematopoietic cells is responsible for the nonspecific immunosuppression that appears to result after blood transfusion in prospective organ transplant recipients (23).

Summary

Graft-vs.-host (GVH)-related immunosuppression has previously been demonstrated in F₁ rodent recipients of parental lymphoid cells, and has been thought to result from an immunologic attack of the donor against the host. Since all cells of such F₁ recipients could potentially bear target class I MHC alloantigens, it has not previously been possible to determine precisely the target tissues responsible for development of GVH-related effects. In the present studies we have used mixed allogeneic chimeras as recipients of host or donor-strain lymphocyte inocula, and have made the surprising observation that "GVH-induced" immune unresponsiveness does not require GVH reactivity, per se, but develops in the presence of a one-way alloresponse against lymphohematopoietic cells in either the GVH or the host-versus-graft direction.

We thank Drs. Gene Shearer and Thor Sundt for helpful review of the manuscript, and Mrs. Judy Jaworek for expert secretarial assistance. We also thank Mr. Donald Holt and his staff for excellent animal care.

Received for publication 4 August 1988.

References

1. Storb, R., and E. D. Thomas. 1985. Graft-versus-host disease in dog and man: the Seattle experience. *Immunol. Rev.* 88:215.
2. Fiscus, W. G., B. T. Morris, J. Session, and J. J. Trentin. 1962. Specificity, host-age effect, and pathology of homologous disease induced in irradiated F₁ hybrid mice by transplantation of parental lymphoid tissue. *Ann. NY Acad. Sci.* 99:355.
3. Holda, J. H., T. Maier, and H. N. Claman. 1985. Murine graft-versus-host disease across minor barriers: immunosuppressive aspects of natural suppressor cells. *Immunol. Rev.* 88:87.
4. Korngold, R., and J. Sprent. 1983. Lethal graft-versus-host disease across minor histocompatibility barriers in mice. *Clin. Haematol.* 12:681.
5. Trentin, J. J. 1956. Mortality and skin transplantability in X-irradiated mice receiving

- isologous, hematologous or heterologous bone marrow. *Proc. Soc. Exp. Biol. Med.* 92:688.
6. Tutschka, P. J., R. Teasdall, W. E. Beschorner, and G. W. Santos. 1982. Chronic graft-versus-host disease in the rat radiation chimera. II. Immunological disease in long-term chimeras. *Transplantation (Baltimore)*. 34:289.
 7. Martinez, C., J. M. Smith, and R. A. Good. 1961. Runt disease in adult tolerant mice induced by intravenous injection of immunologically competent cells. *Proc. Soc. Exp. Biol. Med.* 106:572.
 8. Stastny, P., V. A. Stembridge, and M. Ziff. 1963. Homologous disease in the rat, a model for autoimmune disease. I. General features and cutaneous lesions. *J. Exp. Med.* 118:635.
 9. Lapp, W. S., T. Ghayur, M. Mendes, M. Seddik, and T. A. Seemayer. 1985. The functional and histological basis for graft-versus-host induced immunosuppression. *Immunol. Rev.* 88:107.
 10. Moser, M., T. Iwasaki, and G. M. Shearer. 1985. Cellular interactions in graft-versus-host-induced T cell immune deficiency. *Immunol. Rev.* 88:135.
 11. Moser, M., T. Mizuochi, S. O. Sharrow, A. Singer, and G. M. Shearer. 1987. Graft-vs-host reaction limited to a class II MHC difference results in a selective deficiency in L3T4⁺ but not in Lyt-2⁺ T helper cell function. *J. Immunol.* 138:1355.
 12. Fukuzawa, M., and G. M. Shearer. Graft-vs-host mice exhibit a selective thymus-derived defect in ability to develop MHC self-restricted L3T4 helper cell function. *Immunol. Res.* In press.
 13. Seemayer, T. A., W. S. Lapp, and R. P. Bolande. 1978. Thymic epithelial injury in graft-versus-host reactions following adrenalectomy. *Am. J. Pathol.* 93:325.
 14. Via, C. S., S. O. Sharrow, and G. M. Shearer. 1987. Role of cytotoxic T lymphocytes in the prevention of lupus-like disease occurring in a murine model of graft-vs-host disease. *J. Immunol.* 139:1840.
 15. Billingham, R. E. 1968. The biology of graft-versus-host reactions. *Harvey Lect.* 62:21.
 16. Ildstad, S. T., and D. H. Sachs. 1984. Reconstitution with syngeneic plus allogeneic or xenogeneic bone marrow leads to specific acceptance of allografts or xenografts. *Nature (Lond.)*. 307:168.
 17. Ildstad, S. T., S. M. Wren, S. A. Barbieri, and D. H. Sachs. 1985. Characterization of mixed allogeneic chimeras. Immunocompetence, in vitro reactivity, and genetic specificity of tolerance. *J. Exp. Med.* 162:231.
 18. Ildstad, S. T., S. M. Wren, J. A. Bluestone, S. A. Barbieri, D. Stephany, and D. H. Sachs. 1986. Effect of selective depletion of host and/or donor bone marrow on lymphopoietic repopulation, tolerance, and graft-vs-host disease in mixed allogeneic chimeras (B10 + B10.D2→B10). *J. Immunol.* 136:28.
 19. Sykes, M., A. Eisenthal, and D. H. Sachs. 1988. Mechanism of protection from graft-vs-host disease in murine mixed allogeneic chimeras. I. Development of a null cell population suppressive of cell-mediated lympholysis responses and derived from the syngeneic bone marrow component. *J. Immunol.* 140:2903.
 20. Shearer, G. M., and R. P. Polisson. 1981. Mutual recognition of parental and F1 lymphocytes. III. Parental determinants recognized by F1 host mice in resistance to graft-vs-host-associated immunosuppression map to H-2D^b. *J. Immunol.* 126:545.
 21. Sykes, M., C. H. Chester, and D. H. Sachs. 1988. Protection from graft-versus-host disease in fully allogeneic chimeras by prior administration of T- cell-depleted bone marrow. *Transplantation (Baltimore)*. 46:327.
 22. Sykes, M., M. A. Sheard, and D. H. Sachs. 1988. Effects of T cell depletion in radiation bone marrow chimeras. II. Requirement for allogeneic T cells in the reconstituting bone marrow inoculum for subsequent resistance to breaking of tolerance. *J. Exp. Med.* 168:661.
 23. Klatzmann, D., J. C. Gluckman, F. Chapuis, C. Foucalt, F. Metivier, and J. Rottembourg. 1985. Independent cell-mediated lympholysis and mixed lymphocyte reaction suppression after blood transfusion in humans. *Transplant. Proc.* 17:1051.