

Different Use of T Cell Receptor Transducing Modules in Two Populations of Gut Intraepithelial Lymphocytes Are Related to Distinct Pathways of T Cell Differentiation

By D. Guy-Grand,* B. Rocha,† P. Mintz,* M. Malassis-Seris,* F. Selz,* B. Malissen,§ and P. Vassalli||

From *Institut National de la Santé et de la Recherche Médicale (INSERM) U.132, Hôpital Necker-Enfants Malades, and †U.345, Centre Hospitalo-Universitaire Necker-Enfants Malades, 75015 Paris, France; §INSERM-Centre National de la Recherche Scientifique, Centre d'Immunologie de Marseille-Luminy, 13288 Marseille Cedex 9, France; and ||Department of Pathology, University of Geneva, 1211 Geneva, Switzerland

Summary

Most gut intraepithelial cells (IEL) of the mouse are T cells that bear CD8 molecules, present either as α - β chain heterodimers ($CD8\beta^+$) or as α chain homodimers ($CD8\beta^-$). All $CD8\beta^+$ IEL bear α/β T cell receptors (TCR); $CD8\beta^-$ IEL bear either α/β or γ/δ TCR and are considered to be a thymus-independent (TI) population, probably arising locally from a small fraction of $CD3^-$ IEL containing the recombinant activating gene RAG proteins. Here we report that TI $CD8\beta^-$ IEL, whether bearing α/β or γ/δ TCR, contain, in normal mice, mRNAs for both ζ and Fc ϵ RI γ chains. These chains are present in their $CD3$ -TCR complexes as homo- or heterodimers. In contrast, only ζ chain mRNA and homodimers are found in gut $CD8\alpha/\beta^+$ IEL and in peripheral T lymphocytes. Intestinal $CD3^-$ precursor cells contain only γ chain, and $CD3^-$ IL-2R⁺ thymocyte precursors only ζ chain mRNAs. Only very primitive thymocyte precursors contain detectable γ chain mRNA, and it thus appears that Fc ϵ RI γ chain use is switched off at a very early stage during thymocyte differentiation. Thus, T cell differentiation in the gut epithelium differs from that occurring in the thymus, from which $CD8\beta^+$ IEL appear to derive. Use of different TCR transducing modules and CD8 accessory molecules between the TI and the thymus-derived T cell populations provides an explanation for their difference in reactivity to antigenic stimulations and thus in selection of repertoires.

In the mouse, we have identified two main populations of gut intraepithelial lymphocytes (IEL)¹. One, which bears TCR α/β or γ/δ , is present in significant amounts only in this peculiar localization, and is characterized by the fact that it does not express CD4 and CD8 α/β coreceptors, but CD8 α homodimers ($CD4CD8\beta^-$ IEL) (1). The other, in contrast, bears, as does the bulk of peripheral lymphocytes, TCR α/β and CD4 or CD8 α/β molecules ($CD4CD8\beta^+$ IEL). IEL from both populations are differentiated and functional: they contain granules rich in granzymes and perforin and are cytotoxic (1–3). Since the $CD4CD8\beta^-$ subset develops in the absence of a thymus (1), it is thymus independent (TI), and appears to differentiate locally from $CD3^-$ IEL which contain mRNA for the recombinase activating gene RAG-1 (1, 4). We have presented evidence (1, 5, 6), that, in contrast, the $CD4CD8\beta^+$ IEL are thymus dependent (TD).

Mice mutated in the gene coding for the ζ/η chain of the TCR-CD3 complex have severe defects in thymocyte maturation and in peripheral T cells and gut TD-IEL. In contrast, gut TI-IEL are little affected in these mutant mice, probably because they use Fc ϵ RI γ chain dimers (7) instead of ζ chain dimers (8, 9) as a signal transmitter associated to their TCR-CD3 complex (10–13).

This study analyzes the signal transduction modules present, in normal mice, in the CD3 complexes of the various populations of IEL, peripheral lymphocytes and thymocytes, and explores the presence of mRNAs for ζ and Fc ϵ RI γ chains (referred to below as γ chains) in the precursors of IEL and thymocytes. The resulting observations lead to the conclusion that T cell differentiation in the gut epithelium differs from that occurring in the thymus, and help to delineate two main ontogenic pathways among lymphocytes.

Materials and Methods

Animals. Conventional C3H DBA/2 F₁ mice, C57BL/6 mice transgenic for the α and β chains of a TCR specific for the HY

¹ Abbreviations used in this paper: DN, double negative; IEL, intraepithelial lymphocyte; RAG, recombinant activating gene; TD, thymus dependent; TI, thymus independent; TN, triple negative.

antigen (14) were raised in our animal house. RAG 1^{-/-} (15) and TCR β ^{-/-} (16) mutant mice, a kind gift of P. Mombaerts and S. Tonegawa (Howard Hughes Medical Institute, Massachusetts Institute of Technology, Cambridge, MA 02139), were raised at Centre National de la Recherche Scientifique (Orleans, France), as were RAG-2^{-/-} (17) mutant mice, also used. Swiss nude mice were from Iffa Credo (L'Abresle, France).

Cell Preparations and Separation of Cell Subsets. All subpopulations were sorted with a FACStar[®] flow cytometer (Becton Dickinson & Co., Mountain View, CA) after double labeling with appropriate antibodies. Biotinylated anti-TCR β (H57.597) or δ (GL3) mAbs (revealed with streptavidin-coupled PE), FITC-labeled anti-CD4 (GK1.5) plus CD8 β (H35.17.2) mAbs (1), FITC-labeled PC61 5.3 mAb (CD25) (American Type Culture Collection, Rockville, MD), and biotinylated anti-HY TCR α transgene mAb (18), were used. IEL were isolated as described (19). Peripheral T cells were depleted of B cells by two successive panning on plates coated with an anti-mouse Ig goat antibody. Double negative (DN) TCR⁺ and ⁻ thymocytes were obtained by sorting of cells previously depleted of CD4⁺ CD8⁺ thymocytes by antibody plus complement treatment. In vivo activated T cells were collected from the thoracic duct lymph of F₁ irradiated mice undergoing a graft-vs.-host reaction 5 d after the injection of parental peripheral lymphocytes (20). In vitro activated T cells were obtained by 2-d stimulation of T cells from the thoracic duct lymph with 10 μ g/ml Con A.

Northern Blot Analyses of ζ and Fc ϵ RI γ Chain mRNAs. RNAs were prepared as described (1). The cDNA riboprobes were obtained from full-length cDNAs (21, 22) inserted in pGEM-3 plasmids, and Northern blots were performed in conditions of stringency described previously (1), with hybridization at 62°C and washing at 70°C.

Western Blot Analyses of Anti-CD3 Immunoprecipitates from Cell Lysates. 2–3 \times 10⁶ sorted IEL or LN T cells were solubilized in 1 ml digitonin lysis buffer (1% digitonin, 0.02% N₃Na, 5 mM iodoacetamide, 10 μ g/ml aprotinin, and 100 mM PMSF) for 30 min on ice. Supernatants were precipitated in the digitonin lysis buffer using 5 μ l of protein A-purified hamster anti-CD3 ϵ mAb (2C11) and 50 μ l of a 50% solution of protein A-Sepharose beads (Pharmacia, Uppsala, Sweden). After centrifugation, the beads were incubated in sample buffer (2% SDS, 10% glycerol, 0.08 M Tris-HCl, pH 6.8, 0.02% bromophenol blue), and the supernatant electrophoresed in nonreducing conditions on 12.5% SDS polyacrylamide gels, followed by transfer to nitrocellulose. Coprecipitated chains were revealed with either a rabbit anti-Fc ϵ RI γ antiserum (23) or a rabbit antiserum recognizing a ζ/η peptide (24), followed by incubation with horseradish peroxidase mouse anti-rabbit Ig (The Jackson Laboratory, Bar Harbor, ME) and use of the enhanced chemiluminescence protocol (Amersham International, Amersham, Bucks, UK). Note that for TCR γ/δ ⁺ IEL, lysates from 3 to 4 \times 10⁶ cells were sufficient to clearly observe the protein bands, whereas for TCR α/β ⁺ TI-IEL, lysates from 10 to 12 \times 10⁶ cells, requiring pooling of several cell sorting procedures, were necessary. This difference is probably related to the lower expression of CD3 molecules on TCR α/β ⁺ than on TCR γ/δ ⁺ IEL (data not shown).

Results

Gut IEL Subpopulations Differ in their Use of ζ and γ Chains. Northern blots of mRNA were hybridized with radioactive probes for mRNAs of ζ and γ chains simultaneously, in order to allow a direct comparison of the respective amounts of these mRNAs, identified by their difference in

size. Both mRNAs were found in RNAs from total gut IEL (data not shown). Gut IEL were then fractionated into three populations: CD4CD8 β ⁺ IEL and TCR α/β ⁺ or γ/δ ⁺ CD8 β ⁻ IEL, referred to as TI-IEL (Fig. 1). Whereas ζ chain mRNA was found in all IEL preparations, γ chain mRNA was also present in extracts of the two TI-IEL populations and in higher amounts than ζ chain mRNA. In contrast, no γ chain mRNA was detected in CD4CD8 β ⁺ IEL (Fig. 2). Since a peculiarly small subset of CD4⁺ IEL coexpresses CD8 α chains (probably due to the induction of this last molecule on all subsets of IEL by the gut microenvironment (6, 25), CD4⁺ and CD8 β ⁺ IEL were sorted separately. Both were found to contain only ζ chain transcripts (data not shown). Since TI-IEL can be Thy-1⁺ or ⁻, these subsets were also separately studied. Both contained ζ as well as γ chain mRNAs. TCR⁺ IEL obtained from nude mice (the majority TCR γ/δ ⁺) were also found to contain ζ and γ chain transcripts (Fig. 2). Finally, both mRNAs were also found in the TCR α/β ⁺, CD4CD8 β ⁻ TI-IEL sorted from male mice bearing an anti-HY TCR transgene (Fig. 2) (18).

The presence of ζ and γ chains in the CD3-TCR complexes of TI-IEL was then explored by Western blot analysis of the anti-CD3 precipitates obtained from lysates of TCR α/β ⁺ or γ/δ ⁺ TI-IEL, revealed with anti- ζ/η or γ chain antibodies. Precipitates from lysates of LN cells, used as controls, contained only ζ chain homodimers. In contrast, those from the two subpopulations of TI-IEL, TCR α/β ⁺ or γ/δ ⁺, and those of TCR⁺ IEL from nude mice, contained ζ and γ chain homodimers and ζ/γ heterodimers (Fig. 3). The relative proportions of these various dimers is difficult to evaluate with precision, since it varied somewhat between samples and experiments.

In Various Peripheral T Cell Populations of Euthymic Mice, only ζ Chain mRNA Is Detectable, in Contrast with What Is Observed in Peripheral T Cells of Nude Mice. In euthymic mice, the bulk of lymphocytes from peripheral LN or from the thoracic duct lymph, which are TCR α/β ⁺, CD4⁺, or CD8 β ⁺, contain only ζ chain mRNA (Fig. 4). Peripheral T cells activated in vivo (thoracic duct lymphocytes from mice undergoing an acute graft-vs.-host reaction [20]) or in vitro (2 d of culture) (Fig. 4), which were studied to explore whether




Figure 1. An example of immunofluorescent analysis of the gut IEL used for subpopulation sortings. Only TCR α/β ⁺ IEL bear CD4 or CD8 β coreceptors. The three IEL subpopulations used are easily identified: CD4⁺ or CD8 β ⁺ TCR α/β ⁺ (TD-IEL) (A, quadrant 2 or B, quadrant 4); TI-IEL (CD4⁻ CD8 β ⁻) TCR α/β ⁺ (A, quadrant 1); and TI-IEL (CD4⁻ CD8 β ⁻) TCR γ/δ ⁺ (B, quadrant 1).




Figure 2. Northern blot analyses of ζ and Fc ϵ RI γ chains mRNA in various IEL populations. All Northern blots were hybridized simultaneously with both probes, in order to obtain a quantitative comparison of the presence of the two mRNAs. IEL populations were selected by cell sorting according to their CD4 CD8 β and TCR α/β or γ/δ phenotype as shown in Fig. 1 or: as CD4CD8 β ⁻ Thy1⁻ or ⁺ (TI Thy1⁻ and ⁺), as nude mice TCR⁺ IEL (α/β ⁺ and γ/δ ⁺) (Nude TcR⁺), and as CD8 α/α ⁺ and ⁻ IEL of transgenic male mice bearing the anti-HY TCR transgene (TI TG HY) (18).

short-term T cell activation may induce the expression of γ chains, contained only ζ mRNA. TCR γ/δ ⁺ lymphocytes were obtained by cell sorting of peritoneal cells 3 d after LPS injection (as described [26]): they contained only ζ transcripts (Fig. 4). To study the DN CD4CD8 β ⁻ lymphocytes, which are too scarce in LN from normal mice to be isolated in sufficient number, we selected by cell sorting the DN and CD8 β ^{low} lymphocytes that are present in the peripheral LN of male mice bearing an anti-HY TCR transgene, and the TCR γ/δ ⁺ lymphocytes that represent the only T cell population in TCR β ^{-/-} mutant mice (16). These two populations expressed only ζ chain transcripts (Fig. 4). In contrast, the TCR-CD3⁺ lymphocytes present in the peripheral LN of aged nude mice were found, after sorting, to contain both transcripts (Fig. 4).




Figure 3. Western blot analyses of anti-CD3 immunoprecipitates from cell lysates. In lysates of TCR α/β ⁺ and TCR γ/δ ⁺ CD4CD8 β ⁻ (TI) IEL, ζ homodimers (migrating at 32 Kd), ζ/γ heterodimers (at 24 Kd), and γ homodimers (at 16–18 Kd) are seen, whereas only ζ homodimers are found in lysates of peripheral T lymphocytes. The appearance of γ homodimers as a double band is a common observation (42).




Figure 4. Northern blot analyses of ζ and Fc ϵ RI γ chain mRNAs in various peripheral T cell populations. (TD Lymph) Lymphocytes from the thoracic duct lymph; (GVHR) graft-vs.-host reaction; (Con A) after 2 d in culture with Con A; (IP γ/δ ⁺) intraperitoneal TCR γ/δ ⁺ lymphocytes; (LN TG HY) lymphocytes from LN of transgenic male mice bearing an anti-HY TCR transgene (18) sorted as DN or CD8 α/β ; (LN γ/δ ⁺) TCR γ/δ ⁺ cells obtained from the LN of TCR β ^{-/-} mutant mice (16); (LN Nude TcR⁺) CD3⁺ cells obtained from the LN of nude mice.

Precursors of Thymocytes and of Gut TI-IEL Express γ Homodimers, but in the Thymocyte Differentiation Pathway, Transcription of the γ Chain Is Switched Off Very Early. Precursors of TI-IEL are probably contained in the CD3⁻ IEL population (see Discussion). These cells, sorted from the IEL of normal young mice (since they are in increased percentage compared to adult mice) or of nude mice, contain only γ mRNA (Fig. 5). This is also the case with IEL of RAG^{-/-} mutant mice, which have a block in differentiation at the




Figure 5. Northern blot analyses of IEL precursors and of DN thymocytes. The DN TCR⁺ thymocytes from normal mice, as the DN TCR⁻ cells, contain only ζ chain mRNA, but cannot be subdivided in TCR α/β ⁺ and γ/δ ⁺ cells, since their number is too low. Thus, DN TCR α/β ⁺ thymocytes were obtained by cell sorting from the thymus of male transgenic mice bearing an anti-HY TCR transgene (87% of the thymocytes were DN) and TCR γ/δ ⁺ thymocytes were obtained from β ^{-/-} mutant mice (20% of thymocytes), in smaller number, however, explaining the fainter ζ chain mRNA band.

precursor level (15, 17) and whose IEL are thus all CD3⁻ (Fig. 5).

Since only ζ mRNA is detectable in total thymocytes, subpopulations of thymocytes were studied. CD4⁻CD8⁻ DN thymocytes, sorted in TCR⁻ and TCR⁺ DN cells, contain only ζ chain mRNA (Fig. 5 and data not shown). This last result was confirmed for the two populations of DN TCR⁺ thymocytes, by studying transgenic and mutant TCR β ^{-/-} mice (see above) in which DN TCR α/β ⁺ or γ/δ ⁺ thymocytes can be isolated in sufficient number (Fig. 5). In RAG^{-/-} mutant mice, total thymocytes are present in very reduced number and are all DN, and ζ chain mRNA is abundant but γ chain mRNA is also occasionally detectable in trace amounts. Thus, these thymocytes were sorted into CD25⁻ and CD25⁺ (IL-2R α ⁺) subpopulations, since, in the ontogenic pathway of thymocytes, CD25⁻ lymphocytes are considered the earliest precursors, not yet fully committed to the thymocyte pathway (27). It is striking that whereas little or any γ chain mRNA is detectable in the CD25⁺ thymocytes, both γ and ζ chain mRNAs are found in CD25⁻ thymocytes (Fig. 5).

Discussion

These observations show that the two main populations of gut IEL, in normal adult mice, have different CD3-associated transmission molecules. The first population, which bears TCR α/β and CD4 or CD8 α/β molecules, uses only ζ chains. In this respect, CD4⁺ or CD8 β ⁺ IEL are identical to the CD4⁺ or CD8 β ⁺ TCR α/β ⁺ postthymic cells of peripheral lymphoid organs. This observation is in keeping with the demonstration that all these cells are thymus derived (6). The second population, CD4⁻CD8 β ⁻, TCR α/β ⁺ or γ/δ ⁺, which is TI since its presence does not need that of a thymus, uses, in addition to ζ chains, Fc ϵ RI γ chains, and contain, linked to the CD3 complex, homo- or heterodimeric molecules made of these chains.

To properly evaluate the possible functional and ontogenic meanings of these observations, the various uses or presence of γ and/or ζ chains in T cells in different conditions or at different stages of their ontogenic pathways have to be considered.

It has been observed that activated T cells, in vivo or in culture, may use γ chains in association with, or in place of, ζ chains. The two main populations of IEL have several features of differentiated and activated T cells: they contain cytotoxic granules, display cytotoxicity in redirected lysis assays (2, 3), and bear CD69 (data not shown), a molecule present on activated T cells and on NK cells (28, 29). It does not seem likely that the presence of γ chains in gut TI-IEL merely reflects a recent activation of these cells, since both TD and TI-IEL appear activated whereas only TI-IEL express γ chains. Besides, peripheral T cells activated in vivo (thoracic duct lymphocytes from mice undergoing an acute graft vs. host reaction) or in vitro after 2 d of culture contain only ζ chain mRNA (Fig. 4). However, since at least a fraction of TI-IEL bear α/β TCR known to be potentially autoreactive (30), chronic stimulation of some autoreactive TD-IEL might lead

to the progressive appearance of γ chains within these cells. When a situation of this type is experimentally created, in male transgenic mice bearing an anti-HY TCR transgene, CD8 α/α IEL, which all bear the autoreactive TCR transgene (18), contain γ chain mRNA, but the CD8 α/β ^{low} peripheral T cells bearing the same TCR transgene which are recovered from the LN (having escaped negative selection in the thymus) (31) contain only ζ chain mRNA (Figs. 2 and 4). This observation suggests that it is the site of differentiation of T cells rather than repeated stimulation by (auto)antigens which governs the use of γ chains as CD3-associated transmission molecules.

Whereas all CD4⁺CD8 β ⁺ lymphocytes, whether as IEL or as peripheral lymphocytes, use only ζ chain homodimers, ζ and γ homo- and heterodimers linked to CD3 molecules are found in the IEL of nude mice as in the TI-IEL of euthymic mice. The rare DN TCR α/β ⁺ and γ/δ ⁺ in peripheral organs of euthymic mice contain only ζ chain mRNA, whereas the rare TCR CD3⁺ LN lymphocytes of nude mice contain ζ as well as γ chain mRNA. Thus, the use of ζ chain homodimers is not restricted to lymphocytes of thymic origin. Rather, it appears that it is the use of CD3-transducing molecules with both ζ and γ chains as homo- or heterodimeric molecules which is a hallmark of extrathymic origin, whether in athymic mice or in the TI-IEL of euthymic mice, and whether bearing α/β or γ/δ TCRs². Only studies at the clonal level will allow determination of whether each TI-IEL uses simultaneously these three varieties of transmission molecules or whether their use corresponds to various lymphocyte subpopulations, perhaps at different degrees of maturation.

Comparison of the presence of ζ and γ chain mRNAs in the putative precursors of the TD or TI lineages (the latter being mostly represented, in adult euthymic mice, by TI-IEL) leads to interesting conclusions. The earliest thymic lymphoid cells, which are observed in 14-d fetal thymus, contain γ and ζ chain mRNAs and bear CD16 molecules (35). It has been suggested that these cells represent the earliest thymocyte progenitors and may be common to both T and NK lineages (35, 36). In normal adult thymus, these cells, if present, make up a percentage too minute to be detected, and even after selecting the least differentiated thymocytes, CD4⁻CD8⁻CD3⁻ (triple negative [TN]) cells, γ chain mRNA cannot be detected, whereas ζ chain mRNA is abun-

² The absence of γ chain mRNA in TCR γ/δ ⁺ peripheral lymphocytes (Fig. 4) contrasts with the detection of ζ and γ chains in TCR γ/δ ⁺ cells obtained after culture of splenocytes (32). This may result from the culture conditions in IL-2, since we have also observed the presence of both ζ and γ chain mRNAs after prolonged cultures. Absence of γ chain mRNA in ex vivo conditions may suggest that these peripheral TCR γ/δ ⁺ lymphocytes are thymus derived. The precise nature of the transmission molecules used by the TCR γ/δ ⁺ T cells present in the epithelia of the skin and female reproductive organs, which derive from fetal waves of thymocytes (33), remains to be elucidated. The presence of these cells in ζ ^{-/-} mutant mice has suggested that they might use γ chains (34). It is not known if fetal TCR γ/δ ⁺ thymocytes contain γ chain mRNA, in contrast to adult thymocytes which seem to contain only ζ chain mRNA as shown in the present observation.

dant. Among the thymocytes of $RAG^{-/-}$ mutant mice, chosen to detect more easily early progenitors since thymocyte differentiation is blocked in these mice at an immature TN stage because of their inability to recombine TCR genes, only trace amounts of γ chain mRNA are detectable, whereas ζ chain mRNA is present in large amounts. These $RAG^{-/-}$ thymocytes were sorted into a $CD25^{-}$ (the most undifferentiated) and a $CD25^{+}$ subset (the differentiation stage where the first RAG gene transcripts and the first mRNAs for rearranged β , γ , and δ chains are found in normal mice [37]); both subsets contain ζ chain mRNA, but γ chain mRNA is conspicuously detectable only in $CD25^{-}$ thymocytes. Its presence probably corresponds to a still smaller population of very early progenitors that are $CD16^{+}$, $IL-2R\beta^{+}$ (data not shown) and which has been reported to be found in fetal thymocytes, as mentioned above (38). It thus appears that very early in their differentiation, thymocytes become restricted to ζ chain use, the expression of which is required for further differentiation and maturation of TD lymphocytes. Indeed, in $\zeta/\eta^{-/-}$ mutant mice, thymocyte development is blocked at an early stage, DN thymocytes accumulate, few DP thymocytes are generated, and $CD4\ CD8\beta^{+}$ lymphocytes are scarce in peripheral lymphoid organs and among IEL, and do not express detectable levels of TCR chains (10–13). In conclusion, it appears that, whereas very early thymocyte progenitors contain γ chains, transcription of these chains is very rapidly switched off, so that the differentiation pathway within the thymus eventually rests exclusively on ζ chains, the only chains whose mRNA can be detected in all main maturing populations of thymocytes of normal adult mice, including the DN $TCR\alpha/\beta^{+}$ and γ/δ^{+} thymocytes.

This situation contrasts with that of the pathway of TI-IEL. Although the very early precursors of these cells probably come from the BM or the fetal liver, since lymphocyte lineages may derive from a common ancestor, the more immediate precursors of TI-IEL are probably present in the $CD3^{-}$ IEL. These last cells, whose relative number among IEL is low in euthymic adult mice and higher in young mice before weaning or in nude mice (they are the only IEL found in $RAG^{-/-}$ mice), are characterized by a high proliferation rate and a high level of *c-kit* expression (data not shown), a growth factor receptor mostly expressed on incompletely differentiated hematopoietic cells and on early thymocytes (27). About 20% of them express RAG mRNA (4). All $CD3^{-}$ IEL populations from whatever type of mice, contain γ but not ζ chain mRNA. The expression of ζ chains is not necessary for the development of the ζ and γ chain

containing TI-IEL, since these last cells, in complete contrast with the $CD4\ CD8\beta^{+}$ peripheral lymphocytes and IEL, are present in $\zeta/\eta^{-/-}$ mutant mice, although bearing γ/δ or α/β TCR at lower levels than normal (10). In $\gamma\ Fc\epsilon RI^{-/-}$ mutant mice, thymus development and peripheral T lymphocytes are normal (39). Preliminary studies (Guy-Grand, D., J. V. Ravetch, and P. Vassalli) on IEL have shown that $CD4\ CD8\beta^{+}$ IEL are, as expected, normal, and that TI-IEL are also observed, but bear decreased amounts of α/β or γ/δ TCR. This emphasizes the existence of two pathways of differentiation for T lymphocytes. First, the thymic pathway, using only ζ chain homodimers as $CD3$ -dimeric transmission molecules, except at the earliest stage, does not require expression of γ chains but absolutely requires that of ζ chains. Second, the gut TI pathway, using both γ and ζ chains, is less altered by the lack of either chain. Whether the $CD3$ IEL of $\gamma^{-/-}$ mutant mice substitute ζ chains for the missing γ chains or have some capacity of differentiation without ζ and $\gamma\ Fc\epsilon RI$ chain dimers, since the transducing unit made of $CD3\ \gamma$, δ , and ϵ chains is sufficient to transmit some signals (8), will be established by further study.

In conclusion, epithelial cells of the thymus and of the gut mucosa appear to exert on T lymphocyte precursors, microenvironmental influences inducing distinct ontogenic pathways of T cell differentiation. It is important to recognize that the use of different signal transducing systems by these two distinct pathways is also associated with the use of distinct T cell accessory molecules ($CD8\alpha/\alpha$ instead of $CD8\alpha/\beta$, and lack of $CD28$ [40]). Differences in signal receiving and in signal transducing systems may be equally essential. The thymic pathway, in contrast to the gut IEL pathway, requires the expression of $CD8\beta$ chains, since $CD8\beta$ gene $^{-/-}$ mutant mice have a severe defect in thymocytes and in peripheral T cell differentiation (41). It can be speculated that these differences in signal receiving and transducing machinery are related to essential functional differences in vivo in the TD and TI pathways of differentiation, in particular to contrasting mechanisms of selection in response to antigen during differentiation. Thymocytes undergo an extensive process of antigenic “negative” and “positive” selection, which shapes up the TCR repertoire of peripheral thymic selected T lymphocytes. In contrast, the $CD8\alpha/\alpha$ IEL population does not appear to undergo negative selection (30), and whereas it probably results from a process of selection by antigen (18), this process is different from that of the positive selection seen in the thymus, which probably results from “lower affinity” interactions.

We thank J.-P. Kinet, C. Bonnerot, and S. Nishikawa for advice and generous gift of antibodies; P. Mombaerts, S. Tonegawa, and H. von Boehmer for gifts of mutant mice; Ms. Claude Magnin and Ms. Malika Fontaine for technical assistance; Ms. Jacqueline Ntah for typing the manuscript; and Mr. Jean-Claude Rumbeli and Mr. Etienne Denking for photographic work.

This work was supported by grants from INSERM, Association pour la Recherche sur le Cancer (6662), and from the Swiss National Foundation (31-37516.93).

Address correspondence to Delphine Guy-Grand, INSERM U.132, Hôpital des Enfants Malades, 169 rue de Sèvres, Paris 75763 Cedex 15, France, or to Dr. P. Vassalli, Département de Pathologie, Université de Genève, Faculté de Médecine, 1, rue Michel-Servet, 1211 Genève 4, Switzerland.

Received for publication 29 December 1993 and in revised form 4 April 1994.

References

1. Guy-Grand, D., N. Cerf-Bensussan, B. Malissen, M. Malassis-Seris, C. Briottet, and P. Vassalli. 1991. Two gut intraepithelial CD8⁺ lymphocyte populations with different T cell receptors: a role for the gut epithelium in T cell differentiation. *J. Exp. Med.* 173:471.
2. Guy-Grand, D., M. Malassis-Seris, C. Briottet, and P. Vassalli. 1991. Cytotoxic differentiation of mouse gut thymodependent and independent intraepithelial T lymphocytes is induced locally. Correlation between functional assays, presence of perforin and granzyme transcripts, and cytoplasmic granules. *J. Exp. Med.* 173:1549.
3. Guy-Grand, D., B. Rocha, and P. Vassalli. 1993. Origin and development of gut intraepithelial lymphocytes. In *Mucosal Immunology: Intraepithelial Lymphocytes*. H. Kiyono, editor. Raven Press, New York. 21–31.
4. Guy-Grand, D., C. Vanden Broecke, C. Briottet, M. Malassis-Seris, F. Selz, and P. Vassalli. 1992. Different expression of the recombination activity gene RAG-1 in various populations of thymocytes, peripheral T cells and gut thymus-independent intraepithelial lymphocytes suggests two pathways of T cell receptor rearrangement. *Eur. J. Immunol.* 22:505.
5. Guy-Grand, D., and P. Vassalli. 1993. Gut intraepithelial T lymphocytes. *Curr. Opin. Immunol.* 5:247.
6. Rocha, B., P. Vassalli, and D. Guy-Grand. 1994. Thymic and extrathymic origins of gut intraepithelial lymphocyte populations in mice. *J. Exp. Med.* 180:681.
7. Ravetch, J.V., and J.P. Kinet. 1991. Fc receptors. *Annu. Rev. Immunol.* 9:457.
8. Wegener, A.M.K., F. Letourneur, A. Hoeveler, T. Brocker, F. Luton, and B. Malissen. 1992. The T cell receptor/CD3 complex is composed of at least two autonomous transduction modules. *Cell.* 68:83.
9. Irving, B.A., A.C. Chan, and A. Weiss. 1993. Functional characterization of a signal transducing motif present in the T cell antigen receptor ζ chain. *J. Exp. Med.* 177:1093.
10. Malissen, M., A. Gillet, B. Rocha, J. Trucy, E. Vivier, C. Boyer, F. Köntgen, N. Brun, G. Mazza, E. Spanopoulou et al. 1993. T cell development in mice lacking the CD3- ζ/η gene. *EMBO (Eur. Mol. Biol. Organ.) J.* 12:4347.
11. Ohno, H., T. Aoe, S. Taki, D. Kitamura, Y. Ishida, K. Rajewsky, and T. Saito. 1993. Developmental and functional impairment of T cells in mice lacking CD3 ζ chains. *EMBO (Eur. Mol. Biol. Organ.) J.* 12:4357.
12. Qjian, D., A.I. Sperlino, D.W. Lancki, Y. Tatsumi, T.A. Barrett, J.A. Bluestone, and F.W. Fitch. 1993. The γ chain of the high-affinity receptor for IgE is a major functional subunit of the T-cell antigen receptor complex in $\gamma\delta$ T lymphocytes. *Proc. Natl. Acad. Sci. USA.*
13. Liu, C.P., R. Ueda, J. She, J. Sancho, B. Wang, G. Weddell, J. Loring, C. Kurahara, E.C. Dudley, A. Hayday, et al. 1993. Abnormal T cell development in CD3- $\zeta^{-/-}$ mutant mice and identification of a novel T cell population in the intestine. *EMBO (Eur. Mol. Biol. Organ.) J.* 12:4863.
14. von Boehmer, H. 1990. Developmental biology of T cells in T cell-receptor transgenic mice. *Annu. Rev. Immunol.* 8:531.
15. Mombaerts, P., J. Iacomini, R.S. Johnson, K. Herrup, S. Tonegawa, and V.E. Papaioannou. 1992. RAG-1-deficient mice have no mature B and T lymphocytes. *Cell.* 68:869.
16. Mombaerts, P., A.R. Clarke, M.A. Rudnicki, J. Iacomini, S. Itoharu, J.J. Lafaille, L. Wang, Y. Ichikawa, R. Jaenisch, M.L. Hooper, and S. Tonegawa. 1992. Mutations in T-cell antigen receptor genes α and β block thymocyte development at different stages. *Nature (Lond.)* 360:225.
17. Shinkai, Y., G. Rathbun, K.P. Lam, E.M. Oltz, V. Stewart, M. Mendelsohn, J. Charron, M. Datta, F. Young, A.M. Stall, and F.W. Alt. 1992. RAG-2-deficient mice lack mature lymphocytes owing to inability to initiate V(D)J rearrangement. *Cell.* 68:855.
18. Rocha, B., H. von Boehmer, and D. Guy-Grand. 1992. Selection of intraepithelial lymphocytes with CD8 α/α co-receptors by self-antigen in the murine gut. *Proc. Natl. Acad. Sci. USA.* 89:5336.
19. Guy-Grand, D., C. Griscelli, and P. Vassalli. 1978. The mouse gut lymphocyte, a novel type of T cell. Nature, origin, and traffic in mice in normal and graft-versus-host conditions. *J. Exp. Med.* 148:1661.
20. Guy-Grand, D., and P. Vassalli. 1986. Gut injury in mouse graft-versus-host reaction. Study of its occurrence and mechanisms. *J. Clin. Invest.* 77:1584.
21. Weissman, A.M., M. Baniyash, D. Hou, L.E. Samelson, W.H. Burgess, and R.D. Klausner. 1988. Molecular cloning of the zeta chain of the T cell antigen receptor. *Science (Wash. DC)* 239:1018.
22. Ra, C., M.H.E. Jouvin, and J.P. Kinet. 1989. Complete structure of the mouse mast cell receptor for IgE (FceRI) and surface expression of chimeric receptors (rat-mouse-human) on transfected cells. *J. Biol. Chem.* 264:15323.
23. Letourneur, O., I.C.S. Kennedy, A.T. Brini, J.R. Ortaldo, J.J. O'Shea, and J.P. Kinet. 1991. Characterization of the family of dimers associated with Fc receptors (FceRI and Fc γ RIII). *J. Immunol.* 147:2652.
24. Orloff, D.G., S.J. Frank, F.A. Robey, A.M. Weissman, and R.D. Klausner. 1989. Biochemical characterization of the η chain of the T-cell receptor. A unique subunit related to ζ . *J. Biol. Chem.* 264:14812.
25. Rocha, B., P. Vassalli, and D. Guy-Grand. 1992. The extrathymic T-cell development pathway. *Immunol. Today* 13:449.
26. Skeen, M.J., and H.K. Ziegler. 1993. Intercellular interactions and cytokine responsiveness of peritoneal α/β and γ/δ T cells from *Listeria*-infected mice: synergistic effects of interleukin 1 and 7 on γ/δ T cells. *J. Exp. Med.* 178:985.
27. Godfrey, D.I., and A. Zlotnik. 1993. Control points in early T-cell development. *Immunol. Today* 14:547.
28. Ziegler, S.F., F. Ramsdell, K.A. Hjerrild, R.J. Armitage, K.H. Grabstein, K.B. Hennen, T. Farrar, W.C. Fanslow, E.M. Shevach, and M.R. Alderson. 1993. Molecular characterization of the early activation antigen CD69: a type II membrane glycoprotein related to a family of natural killer cell activation

- antigens. *Eur. J. Immunol.* 23:1643.
29. López-Cabrera, M., A.G. Santis, E. Fernández-Ruiz, R. Blacher, F. Esch, P. Sánchez-Mateos, and F. Sánchez-Madrid. 1993. Molecular cloning, expression, and chromosomal localization of the human earliest lymphocyte activation antigen AIM/CD69, a new member of the C-type animal lectin superfamily of signal-transmitting receptors. *J. Exp. Med.* 178:537.
 30. Rocha, B., P. Vassalli, and D. Guy-Grand. 1991. The V β repertoire of mouse gut homodimeric α CD8⁺ intraepithelial T cell receptor α/β ⁺ lymphocytes reveals a major extrathymic pathway of T cell differentiation. *J. Exp. Med.* 173:483.
 31. Rocha, B., C. Tanchot, and H. von Boehmer. 1993. Clonal anergy blocks in vivo growth of mature T cells and can be reversed in the absence of antigen. *J. Exp. Med.* 177:1517.
 32. Qian, D., A.I. Sperling, D.W. Lancki, Y. Tatum, T.A. Berrett, J.A. Bluestone, and F.W. Finch. 1993. The γ chain of high-affinity receptor for IgE is a major functional subunit of the T-cell antigen receptor complex in $\gamma\delta$ T lymphocytes. *Proc. Natl. Acad. Sci. USA.* 90:11875.
 33. Lafaille, J.J., A. DeCloux, M. Bonneville, Y. Takagaki, and S. Tonegawa. 1989. Junctional sequences of T cell receptor $\gamma\delta$ genes: implications for $\gamma\delta$ T cell lineages and for a novel intermediate of V-(D)-J joining. T cell 46E1. *Cell.* 59:859.
 34. Ohno, H., S. Ono, N. Hirayama, S. Shimada, and T. Saito. 1994. Preferential usage of the Fc receptor γ chain in the T cell antigen receptor complex by γ/δ T cells localized in epithelia. *J. Exp. Med.* 179:365.
 35. Rodewald, H.R., P. Moingeon, J.L. Lucich, C. Dosiou, P. Lopez, and E.L. Reinherz. 1992. A population of early fetal thymocytes expressing Fc γ RII/III contains precursors of T lymphocytes and natural killer cells. *Cell.* 69:139.
 36. Moingeon, P., H.R. Rodewald, D. McConkey, A. Mildonian, K. Awad, and E.L. Reinherz. 1993. Generation of natural killer cells from both Fc γ RII/III⁺ and Fc γ RII/III⁻ murine fetal liver progenitors. *Blood.* 82:1453.
 37. Wilson, A., W. Held, and H.R. Macdonald. 1994. Two waves of recombinase gene expression in developing thymocytes. *J. Exp. Med.* 179:1355.
 38. Falk, I., C.N. Levelt, and K. Eichmann. 1993. Lineage relationships of the fetal thymocyte subset that expresses the β chain of the interleukin-2 receptor. *Eur. J. Immunol.* 23:3373.
 39. Takai, T., M. Li, D. Sylvestre, R. Clynes, and J.V. Ravetch. 1994. FcR γ chain deletion results in pleiotropic effector cell defects. *Cell.* 76:519.
 40. Ohteki, T., and H.R. Macdonald. 1993. Expression of the CD28 costimulatory molecule on subsets of murine intestinal intraepithelial lymphocytes correlates with lineage and responsiveness. *Eur. J. Immunol.* 23:1251.
 41. Nakayama, K.I., K. Nakayama, I. Negishi, K. Kuida, M.C. Louie, O. Kanagawa, H. Nakauchi, and D.Y. Loh. 1994. Requirement for CD8 β chain in positive selection of CD8-lineage T cells. *Science (Wash. DC).* 263:1131.
 42. Paolini, R., and J.P. Kinet. 1993. Cell surface control of the multiubiquitination and deubiquitination of high-affinity immunoglobulin E receptors. *EMBO (Eur. Mol. Biol. Organ.) J.* 12:779.