

Very Low Affinity B Cells Form Germinal Centers, Become Memory B Cells, and Participate in Secondary Immune Responses When Higher Affinity Competition Is Reduced

Joseph M. Dal Porto,¹ Ann M. Haberman,² Garnett Kelsoe,³
and Mark J. Shlomchik²

¹Department of Microbiology and Immunology, University of Maryland School of Medicine, Baltimore, MD 21201

²Department of Laboratory Medicine and Section of Immunobiology, Yale University School of Medicine, New Haven, CT 06510

³Department of Immunology, Duke University Medical Center, Durham, NC 27710

Abstract

To understand the relationship between the affinity of the B cell antigen receptor (BCR) and the immune response to antigen, two lines of immunoglobulin H chain transgenic (Tg) mice were created. H50G μ^a and T1(V23) μ^a mice express μ H chain transgenes that associate with the λ 1 L chains to bind the (4-hydroxy-3-nitrophenyl)acetyl hapten with association constants (K_s) of only $1.2 \times 10^5 \text{ M}^{-1}$ and $3 \times 10^4 \text{ M}^{-1}$, respectively. Both lines mounted substantial antibody-forming cell (AFC) and germinal center (GC) responses. H50G μ^a Tg mice also generated memory B cells. T1(V23) μ^a B cells formed AFC and GCs, but were largely replaced in late GCs by antigen-specific cells that express endogenous BCRs. Thus, B lymphocytes carrying BCRs with affinities previously thought to be irrelevant in specific immune responses are in fact capable of complete T cell-dependent immune responses when relieved of substantial competition from other B cells. The failure to observe such B cells normally in late primary responses and in memory B cell populations is the result of competition, rather than an intrinsic inability of low affinity B cells.

Key words: mutation • antibody affinity • plasma cell • variable region gene • λ 1 immunoglobulin

Introduction

Early humoral responses are heterogeneous and mainly low affinity (1–4). As the response progresses, Ab diversity is reduced and affinity increases (5–7). It is possible that early low affinity Ab is biologically irrelevant and the result of nonspecific or bystander stimulation (8, 9). Alternatively, very low affinity B cells may respond specifically but are lost as the response progresses. For example, intrinsic affinity requirements could limit cell participation in humoral responses. Survival in germinal centers (GCs) or memory differentiation may require signaling above a threshold (10–12). Alternatively, the loss of low affinity B cells from

immune responses could reflect competition among antigen-reactive B cells (13).

To distinguish these models for the antigen-driven selection of B cells, we generated the following two lines of transgenic (Tg) mice expressing Ig H chain Tgs that encode extremely low affinity anti-(4-hydroxy-3-nitrophenyl)acetyl (NP) antibodies when associated with the λ 1 L chain: $1.2 \times 10^5 \text{ M}^{-1}$ for H50G μ^a and $\sim 3.0 \times 10^4 \text{ M}^{-1}$ for T1(V23) μ^a .

Materials and Methods

Mice. 6–8-wk-old C.B-17 and C.B-17-SCID mice were purchased from Taconic Farms, Inc. All mice were maintained in sterile microisolator cages.

Creation of H50G μ^a and T1(V23) μ^a Tg Mice. The VH186.2/DFL16.1 rearrangement, containing a single mutation at codon 50, designated H50G and V23 analogue VH rearrangement from an unmutated VDJ rearrangement of the V23 gene taken from a C57BL/6 mouse 10 d after immunization with NP, were subcloned into a vector that contains the 11.6-kb C μ fragment that

The online version of this article contains supplemental material.

G. Kelsoe and M.J. Shlomchik contributed equally to this work.

Address correspondence to Mark J. Shlomchik, Department of Laboratory Medicine and Section of Immunobiology, Yale University School of Medicine, 330 Cedar St. Rm CB465, New Haven, CT 06510. Phone: 203-688-2089; Fax: 203-688-2748; E-mail: mark.shlomchik@yale.edu; or Garnett Kelsoe, Department of Immunology, Duke University Medical Center, 117 Jones Bldg., DUMC 3010, Research Dr., Durham, NC 27710. Phone: 919-613-7815; Fax: 919-613-7878; E-mail: ghkelsoe@duke.edu

includes the membrane and secretory exons (14). The H50G μ^a lineage had two copies of the Tg, whereas two separate T1(V23) μ^a lineages with similar phenotypes were used with two to three copies or a single copy of the Tg. Tg mice were backcrossed with C.B-17 mice for five or more generations.

Flow Cytometry. The following antibodies were used: phycoerythrin-conjugated anti-B220 (CD45RA-PE; BD PharMingen); fluorescein-conjugated anti-mouse IgM^a (RS3.1-FITC); biotinylated AF6-7.8 (anti-IgM^b); fluorescein-conjugated or biotinylated Ls136 (anti- λ 1); and biotinylated goat anti-mouse κ L chain-specific and biotinylated goat anti-mouse Ig (Southern Biotechnology Associates, Inc.). Red 670-conjugated streptavidin (GIBCO BRL) was used to reveal biotin-coupled Ab staining.

Antigens and Immunizations. Mice were immunized intraperitoneally with 50 μ g of NP₁₆-chicken γ -globulin (CG), (4-hydroxy-5-iodo-3-nitrophenyl)acetyl (NIP)₁₂-CG, or CG precipitated in alum. For secondary/memory responses and challenge of repopulated SCID mice, 20 μ g of soluble NP₁₆-CG in PBS was administered intravenously on the indicated days.

Immunohistology. Immunohistology was performed as previously described (3, 15). GCs were enumerated as peanut agglutinin (PNA)⁺ areas within lymphoid follicles (\pm SD from three separate sections). Endogenous Ig-expressing B cells in the GC of H50G μ^a or T1(V23) μ^a mice were enumerated using both AF6-7.8-biotin and biotinylated goat anti-mouse IgG₁, followed by streptavidin-alkaline phosphatase. To detect apoptotic cells, TUNEL assays were performed using in situ apoptosis detection kits (Oncor).

Microdissection of Cells, DNA Amplification, and Sequencing. Microdissection of cells, DNA amplification, and sequencing was performed as previously described (14, 16). Cellular material (constituting 10–20 cells) was microdissected from individual GCs (3, 15, 17). In H50G μ^a mice, GCs sampled were also IgM^{a+} and devoid of endogenous Ig-expressing (IgM^{b+} or IgG₁⁺) cells. DNA was amplified by Pfu polymerase (Stratagene) using nested primers specific for V λ 1 and J λ 1 elements (14).

Quantitation of Serum Ab. Ab titers were determined by endpoint dilution in ELISA using plates coated with NP₂₁-BSA, NIP₁₉-BSA, or CG (7). The endpoint was the last serial dilution that demonstrated a signal greater than twofold above the background. NP-specific ELISAs using differentially haptenated substrates were used to quantitate high and low affinity Ab (3).

Reduction of Serum Ab by 2-Mercaptoethanol (2-ME). Serum was assessed for NP reactivity by ELISA after treatment with 2-ME (3). For calculations of the percent of the IgM^a titer resistant to 2-ME treatment, mock and 2-ME ELISA results were converted to relative Ab concentrations by comparison to a standard purified H50G μ^a / λ 1 transfectoma Ab.

Adoptive Transfers of Splenocytes. 5×10^6 splenocytes recovered from H50G μ^a and C.B-17 mice 30 d after immunization with 50 μ g of NP₁₂-CG precipitated in alum (18) were injected intravenously into recipient SCID mice. Recipients were challenged with 20 μ g of NP₁₂-CG in PBS or PBS alone 24 h later. 9 d after immunization the recipient sera were collected.

Online Supplemental Material. The supplementary material shows original V λ sequence data as well as summary tables that analyze the distribution of mutations in the sequences. Fig. S1 shows all codons that contained mutations in V λ 1 sequences derived from either C.B-17 or H50G μ^a mice, as described in the Materials and Methods and Results sections. Table S1 is a comparison of the two datasets, showing the frequency of overall mutations as well as replacement and silent mutations. Table S2 shows the ratios of replacement and silent ratios in each subregion

of V λ . Online supplemental material available at <http://www.jem.org/cgi/content/full/jem.20011550/CD1>.

Results

Characterization of H50G μ^a and T1(V23) μ^a Tg Mice. Splenic B cells from naive H50G μ^a and T1(V23) μ^a mice demonstrated efficient allelic exclusion, with $\leq 2\%$ expressing endogenous IgM^b protein (unpublished data). The ratio of λ 1: κ L chain-bearing B cells were comparable in both Tg and control mice, and all Tg lines displayed total surface Ig densities comparable to normal, naive B cells (unpublished data).

Humoral Immune Responses in H50G μ^a Tg Mice. Naive H50G μ^a Tg mice had low titers ($\leq 1:60$) of NP-specific serum Ab that sharply increased after immunization with NP-CG (Fig. 1 A). At day 12, the titers of NP-binding Tg (IgM^a) Ab increased 1,000-fold in H50G μ^a mice. Only in 3 out of 31 H50G μ^a mice were substantial levels of IgG₁ anti-NP detected (Fig. 1 A), indicating that typically most anti-NP was Tg derived. Levels for anti-NP λ 1 and total Ig at day four were similar for Tg and control animals. However, C.B-17 mice later achieved 5–10-fold higher titers. To confirm that H50G μ^a mice expressed only low affinity Tg-encoded antibodies in response to immunization, sera of H50G μ^a and C.B-17 mice were mildly reduced with 2-ME (3). The binding of very low affinity IgM Ab is avidity dependent and lost upon 2-ME reduction (5), whereas higher affinity IgM or IgG Abs are unaffected. The reduction of Tg sera led to a near complete loss ($\geq 97\%$) of NP reactivity (Fig. 1 B). This loss was comparable to that observed for purified H50G μ^a / λ 1 transfectoma Ab (3). In contrast, C.B-17 serum titers were unaffected.

Immune Response of Very Low Affinity B Cells. To directly test whether B cells with very low affinities were capable of initiating and sustaining specific humoral responses, H50G μ^a Tg mice were immunized with NIP-CG and T1(V23) μ^a Tg mice were challenged with NP-CG and NIP-CG. H50G μ^a / λ 1 Ab has a lower affinity for NIP (6.3×10^4 M⁻¹) than for NP (3). T1(V23) μ^a / λ 1 Ab has a $K_a = 5.0 \times 10^4$ M⁻¹ for NIP and an affinity for NP that is too low for reliable measurement by fluorescence quenching (5).

T1(V23) μ^a mice had 8–10-fold increases in hapten-specific IgM^a Ab between 4 and 16 d after immunization with either immunogen, with some mice showing 20–30-fold increases in Ab titer (Fig. 1, C and D; \blacktriangle , \triangle). IgM^a responses by T1(V23) μ^a mice to NP-CG or NIP-CG were lower (5–10-fold) than those observed in H50G μ^a mice for the same immunogen (Fig. 1, C and D; \blacktriangle , \triangle vs. \circ). IgG₁ responses, presumably encoded by the endogenous *Igh* loci, were higher in T1(V23) μ^a mice compared with H50G μ^a mice. The higher levels of endogenously derived anti-NP antibodies in T1(V23) μ^a mice may reflect the extraordinarily poor affinity of these Tg B cells, which confer little advantage in competition with rare B cells that express endogenous VH genes. B cells expressing endogenously encoded VH genes comprise 1–3% of the initial repertoire,




Figure 1. Humoral immune response in low affinity H50G μ^3 and T1(V23) μ^3 Tg mice. (A) H50G μ^3 (○) and C.B-17 controls (●) were injected intraperitoneally with NP-CG in alum. Serial dilutions of sera were assayed for NP binding of total Ig, Tg IgM 3 , anti-NP characteristic λ 1, and IgG $_1$. *, H50G μ^3 mice in which significant anti-NP IgG $_1$ titers may have affected the measurement of IgM 3 Tg levels. (B) Sera from H50G μ^3 and control C.B-17 mice after immunization with NP-CG were assayed for their ability to bind NP-BSA after 2-ME treatment (shaded bars) or mock treatment (solid bars). The change in the relative NP binding of treated samples is shown as a percentage change from mock-treated samples. Values represent the mean values (\pm SD) obtained from three independent experiments. T1(V23) μ^3 Tg mice (\blacktriangle) were compared with H50G μ^3 (○) and C.B-17 (●) mice in response to (C) NP-CG or (D) NIP-CG in alum. Sera were assayed for NP-binding components. Each symbol represents the average endpoint dilution value of serum from an individual mouse measured in triplicate.

of which only a small fraction of B cells would have NP/NIP specificity. Thus, the emergence of these B cells in response to NP/NIP reflects the competition between rare, presumably higher affinity B cells and the much more common low affinity Tg B cells.

At the height of the response, H50G μ^3 mice challenged with NIP had a 100-fold increase in hapten-binding IgM 3 Ab titers compared with controls (Fig. 1 D), which is 2–10-fold lower than those after challenge with NP-CG (Fig. 1, C and D). This observation establishes a relationship between B cell antigen receptor (BCR) affinity/avidity and the magnitude of the Ab response in a single Tg mouse line.

Low Affinity Tg B Cells Can Initiate Primary GCs. PNA $^+$ GCs in H50G μ^3 mice 12 d after immunization (Fig. 2 A) were observed in numbers (\sim 70–80 per section) equivalent to normal mice (6, 15). $65 \pm 5.8\%$ of the GCs in H50G μ^3 mice were stained with both anti-IgM 3 and anti- λ 1 Ab, indicating that they contained NP-specific B cells expressing the H50G μ^3 Tg (Fig. 2, A and C). The percentage of λ 1 $^+$ GCs was only slightly higher than that seen in non-Tg C.B-17 mice for the same day, which was

$54 \pm 3.4\%$. Many B cells in the splenic red pulp exhibited strong cytoplasmic staining with λ 1- and IgM 3 -specific Ab (unpublished data), the characteristic phenotype of plasmablasts and antibody-forming cell (AFC) (18).

H50G μ^3 Tg B cells also formed Tg $^+$ GCs when their BCR affinity was less than 10^5 M $^{-1}$, i.e., after immunization with NIP. Such mice had PNA $^+$ and IgM 3 GCs in numbers equivalent to C.B-17 controls, although many of these GCs were smaller than those in control animals (unpublished data). We also investigated the capacity of very low affinity B cells to form GC in T1(V23) μ^3 mice. Surprisingly, there was no difference in GC number and size in T1(V23) μ^3 mice challenged with NP or NIP compared with C.B-17 controls. However, \sim 80% of GCs in T1(V23) μ^3 mice contained both Tg IgM 3 - and endogenous Ig-expressing B cells (unpublished data). When present, endogenous Ig-expressing B cells (demonstrated with anti-IgM 3 Ab and anti-IgG $_1$ Ab) constituted as much as 50% of the PNA $^+$ cells in each GC. Presumably these endogenously derived B cells reflect the competitive success of rare, higher affinity B cells. To confirm this, we microdissected and sequenced several clusters of λ 1 $^+$ B cells




Figure 2. GC formation in H50G μ^3 mice 12 d after immunization with NP-CG. Splenic serial sections reveal PNA⁺ GC B cells (red) that were labeled in tandem (blue) for: (A) Tg IgM^{a+}; (B) endogenous IgM^{b+}; (C) λ 1 L chain; and (D) κ L chain. rp, red pulp; pals, periarteriolar lymphoid sheath; gc, germinal center. Note that surface Ig staining is weak in GCs due to marked surface IgM down-regulation in GCs. However, the difference between the (A) IgM^a and (B) IgM^b staining of the same GC can be appreciated by how the IgM^a staining modifies the color of the PNA-stained GC cells. A similar picture is seen comparing (C) λ 1 and (D) κ staining. $\times 100$.

costaining for IgM^b/IgG₁ and sequenced their VH regions. In fact, >75% of recovered sequences were V_H186.2, the canonical VH of anti-NP responses, which established that the endogenously derived B cells are higher affinity anti-NP B cells. It is likely that the population of B cells expressing endogenous VDJ rearrangements accounts for the normal size and frequencies of GCs in immunized T1(V23) μ^a animals.

Late GCs of H50G μ^a Mice Show Evidence of Increased Apoptosis. 12 d after immunization with NP-CG, B cells bearing BCRs encoded by the endogenous *Igh^b* loci are infrequent in the GC of H50G μ^a Tg mice (Fig. 3 A). However, at day 16 of the primary response, endogenous Ig-expressing B cells (IgM^{b+} or IgG₁⁺) begin to appear in the GCs, increasing to 5–10% of all PNA⁺ GC cells by day 20 (Fig. 3, B and C). Late GCs (16–20 d after immunization) of H50G μ^a mice had two- to threefold more apoptotic B cells compared with C.B-17 controls, as indicated by TUNEL labeling (Fig. 3, D and E). GCs ($n = 30$) sampled from six H50G μ^a mice at 16 or 20 d after immunization had an average 32% ($\pm 10\%$) TUNEL⁺/PNA⁺/IgD⁻ cells. This was in contrast to GCs ($n = 20$) from four C.B-17 mice at similar days, which contained only 12% ($\pm 7\%$) TUNEL⁺ GC cells.

Somatic Mutation in H50G μ^a Mice. Somatic hypermutation was assayed in the endogenous λ 1 L chain gene of H50G μ^a mice, as ectopic heavy chain Tg generally does not mutate. IgM^{a+}/ λ 1⁺, GC cells from three H50G μ^a mice, and three C.B-17 mice were microdissected and their V λ regions amplified and sequenced (Fig. S1 and Tables S1 and S2). Of 28 H50G μ^a V λ 1/J λ 1 rearrangements




Figure 3. GCs in H50G μ^a mice have increased frequencies of endogenous BCRs and enhanced apoptosis. PNA⁺ GCs (red) from H50G μ^a mice (A) 12, (B) 16, (C) and 20 d after immunization with NP-CG were stained for the presence of endogenous Ig (blue, IgM^b or IgG₁). For detecting apoptotic GC cells, (D) H50G μ^a and (E) C.B-17 sections of spleen from mice at day 16 of the primary immune response were labeled by TUNEL⁺ (red) and with anti-IgM (blue). GCs appear as faint blue areas within darker blue follicles due to surface Ig downmodulation on GC B cells.

examined, 22 were mutated with an average of 5.1 (± 3.7) mutations per rearrangement.

The distribution and type of somatic mutations observed in V λ 1 genes present in Tg and control GC B cells is summarized in Tables SI and SII. 57 of 68 H50G μ^a V λ 1 mutations encoded amino acid replacements. The 68 unique mutations were a striking contrast to the 11 unique mutations found in C.B-17 GCs. Mutations in the H50G μ^a V λ 1/J λ 1 rearrangements had an overall replacement/silent (R/S) ratio three times greater than C.B-17 rearrangements. In addition, the R/S ratio of mutations in CDR1 and CDR2 of H50G μ^a V λ 1 genes was 10- and 6-fold higher, respectively, than the predicted ratios based on codon usage. These data demonstrate enhanced mutation and selection in the endogenous V λ 1 gene rearrangements of H50G μ^a GC B cells.

Low Affinity Primary and Memory Responses in H50G μ^a Mice. To determine the degree of affinity maturation in the primary and memory responses of H50G μ^a mice, Tg and C.B-17 mice were immunized with NP-CG and 60 d later were rechallenged intravenously with soluble NP-CG. Fig. 4 shows the change in the NP-specific IgM^a-Tg Ab titers. Upon secondary immunization, a prompt and significant increase in NP-specific IgM^a Ab was observed. By day 9 of the secondary response, Tg-IgM^a endpoint titers increased from 1:4,500 to 1:72,000 (Fig. 4, \circ). No significant Ab response (>1 out of 200) was observed in unprimed H50G μ^a mice given the same soluble antigen (Fig. 4, ∇).

Affinity maturation, presumably through the mutation of V λ 1, did enhance the avidity of H50G μ^a Ab, as re-




Figure 4. Affinity maturation of the humoral response in H50G μ^3 mice. Bars show the NP5/NP22 binding ratio for IgM 3 Ab. Endpoint titers for NP $_5$ - (Δ) and NP $_{22}$ - (\circ) binding IgM 3 from H50G μ^3 mice are superimposed to demonstrate the actual Ab concentration shifts during the response. The response of naive H50G μ^3 mice to a primary intravenous immunization with soluble NP-CG is also shown (∇). Values represent the mean \pm SD (error bars) of four or five mice for each time point.

flected by the increased portion that bound NP $_5$ -BSA (Fig. 4; Δ , bars). However, this affinity maturation was modest, as 85–90% of secondary Ab generated by H50G μ^3 mice remained sensitive to 2-ME treatment (unpublished data). Interestingly, the rechallenge of primed H50G μ^3 mice with soluble NP-CG elicited a rapid increase in low affinity Ab, as inferred from the precipitous decrease in the NP $_5$ /NP $_{22}$ binding ratio of the serum Ab (Fig. 4, bars). Only 30% of the NP-specific IgM 3 titer at day 9 of the secondary response showed the ability to bind NP $_5$ -BSA. This is a contrast to the 70% observed in the late primary response (Fig. 4) and consistent with the reactivation of low affinity memory B cells and their differentiation into AFC (7).

To exclude that T cell priming alone was responsible for the memory response of H50G μ^3 mice, Tg and control mice were primed with the carrier protein CG, and 60 d later challenged intravenously with soluble NP-CG. Average NP-specific $\lambda 1^+$ Ab titers from CG-primed H50G μ^3 mice were 15-fold lower than cohorts primed with NP-CG (Table I; 1:2,200 vs. 1:33,400). On challenge with soluble antigen, NP-CG-primed H50G μ^3 mice exhibited a 38% absolute increase in the serum Ab titer capable of binding NP $_5$ -BSA, whereas there was no appreciable increase in the NP $_5$ -BSA-binding titer in CG-primed H50G μ^3 mice. Thus, T cell priming alone cannot account for the extent of low affinity Ab elicited in the memory responses of H50G μ^3 mice.

To minimize the influence of preformed Ab on secondary responses, splenocytes from naive or NP-CG-primed (day 30) H50G μ^3 and C.B-17 mice were adoptively transferred into SCID recipients, which were then challenged intravenously with soluble NP-CG. SCID mice that received primed H50G μ^3 splenocytes generated Ab titers that were more than 100-fold higher than SCIDs that were repopulated with naive H50G μ^3 splenocytes. The magnitude of the increase was similar in control mice receiving C.B-17 donor cells (unpublished data). These differences in serum Ab were not due to the transfer of residual AFC, as SCID mice that received primed H50G μ^3 splenocytes but were not challenged with antigen had no appreciable anti-NP serum Ab (titer of 1:60).

Discussion

Normally, very low affinity cells are rare in late GCs, in memory cell populations, or as long-lived AFC (19, 20). By fixing the B cell repertoire at low affinity using a non-mutating ectopic Tg, we show that the virtual absence of low affinity cells in these compartments is due to competition from higher affinity cells. When clonal competition is reduced, very low affinity B cells fill GCs and contribute to memory responses.

Although competition is reduced, low numbers of B cells expressing endogenous VH gene segments provide detectable interclonal competition. Endogenously derived cells are rare in H50G μ^3 Tg mice, especially early in the response. However, many B cells expressing non-Tg BCR emerge in immunized T1(V23) μ^3 mice, and to a lesser ex-

Table I. Memory Response to Soluble Antigen from NP-CG Primed and Carrier Primed Mice^a

Immunization Protocol ^b		Serum Response ^c (end-point dilution)		
Mice	1 ^o (i.p.)	1 ^o (i.v.)	Percent NP5/NP22 Ratio	
CB-17 H50G μ^3	NP-CG	NP-CG	135,826 \pm 69,607	53
		CG	33,417 \pm 10,919	38
C.B-17 H50G μ^3	CG	NP-CG	4,797 \pm 2,010	23
		CG	2,218 \pm 1,581	8
C.B-17 H50G μ^3	none	NP-CG	130 \pm 60	0
		NP-CG	60 \pm 20	0

^aSerum data were obtained from mice 9 d after secondary challenge with soluble antigen.

^bPrimary immunizations were by intraperitoneal administration of 50 μ g NP-CG or CG alone in alum. Secondary/memory responses were elicited through the intravenous injection of 20 μ g soluble NP-CG 60 d after primary challenge.

^cThe NP-specific $\lambda 1^+$ end-point serum titer was used as a measure of the secondary immune response to allow direct comparison between H50G μ^3 and C.B-17 mice.

tent H50G μ^a mice, as demonstrated by the presence of non-Tg Ab (Fig. 1) and histology (Fig. 3). Presumably, the extremely low affinity of T1(V23) μ^a B cells for the NP and NIP haptens allows them to be overwhelmed by rare B lymphocytes that escape allelic exclusion and respond to NP/NIP, as indicated by their frequent expression of V_H186.2, a hallmark of high affinity NP Ab. Thus, higher affinity B cells expressing endogenous genes outcompete a numerous but very low affinity Tg population.

As V(D)J hypermutation creates L chain diversity in GCs (Figure S1), intraclonal competition also exists. R/S ratios in V λ CDR1 and V λ CDR2 suggest strong selection for GC B cell mutants. Although V λ mutations can improve BCR affinity in H50G μ^a mice, the low affinity of secondary Ab in these mice (Fig. 4) suggests that this pathway for affinity maturation is limited. Nonetheless, the high frequency (1.4%) of point mutations in the V λ 1 rearrangements of H50G μ^a GC B cells is almost twice that found in a similar mouse with a germline affinity, V_H 186.2 Tg (14), and much higher than wild-type mice. The high frequency and intense selection of V λ 1 mutations in H50G μ^a mice could reflect the difficulty in improving BCR affinity through L chain mutations alone. However, mutation rates may be intrinsically higher in low affinity B cells.

Our data do not rule out the role(s) for differential signaling governed by affinity/avidity in determining B cell fates (11, 12). Aside from memory development, other aspects of B cell differentiation like isotype switching could be governed by BCR signal strength.

The ability of very low affinity B cells to establish GCs and humoral memory has implications for understanding natural B cell responses. Low affinity BCRs can acquire higher affinities with a single point mutation (19, 21), and the many low affinity B cells that reach GCs are the potential precursors of effective memory B cells. This view is also consistent with direct estimates of clonal diversity in nascent GCs (6) and computer models, which suggest that GCs are seeded by 40–50 precursors (22). Finally, the latent genetic potential of low affinity clones could underlie “clonal shifts” in secondary responses whereby dominant, primary clonotypes are replaced by mutated, dissimilar clones (2, 23).

This work was supported by National Institutes of Health grants AI24335, AG10207, and AG13789 to G. Kelsoe, and AI43603 to M.J. Shlomchik. A.M. Haberman was the recipient of a Donaghue Foundation postdoctoral fellowship.

Submitted: 10 September 2001

Revised: 8 March 2002

Accepted: 13 March 2002

References

1. Chua, M.M., S.H. Goodgal, and F. Karush. 1987. Germ-line affinity and germ-line variable-region genes in the B cell response. *J. Immunol.* 138:1281–1288.
2. Clarke, S.H., L.M. Staudt, J. Kavaler, D. Schwartz, W.U. Gerhard, and M.G. Weigert. 1990. V region gene usage and somatic mutation in the primary and secondary responses to influenza virus hemagglutinin. *J. Immunol.* 144:2795–2801.
3. Dal Porto, J.M., A.M. Haberman, M.J. Shlomchik, and G. Kelsoe. 1998. Antigen drives very low affinity B cells to become plasmacytes and enter germinal centers. *J. Immunol.* 161:5373–5381.
4. Eisen, H.N., and G.W. Siskind. 1964. Variations in affinity of antibodies during the immune response. *Biochemistry.* 3:996–999.
5. Berek, C., and M. Ziegner. 1993. The maturation of the immune response. *Immunol. Today.* 14:400–404.
6. Jacob, J., J. Przylepa, C. Miller, and G. Kelsoe. 1993. In situ studies of the primary immune response to (4-hydroxy-3-nitrophenyl)acetyl. III. The kinetics of V region mutation and selection in germinal center B cells. *J. Exp. Med.* 178:1293–1307.
7. Takahashi, Y., P.R. Dutta, D.M. Cerasoli, and G. Kelsoe. 1998. In situ studies of the primary immune response to (4-hydroxy-3-nitrophenyl)acetyl. V. Affinity maturation develops in two stages of clonal selection. *J. Exp. Med.* 187:885–895.
8. Conger, J.D., B.L. Pike, and G.J. Nossal. 1988. Analysis of the B lymphocyte repertoire by polyclonal activation. Hindrance by clones yielding antibodies which bind promiscuously to plastic. *J. Immunol. Methods.* 106:181–189.
9. Urbain-Vansanten, G. 1970. Concomitant synthesis, in separate cells, of non-reactive immunoglobulins and specific antibodies after immunization with tobacco mosaic virus. *Immunology.* 19:783–797.
10. Madrenas, J., R.L. Wange, J.L. Wang, N. Isakov, L.E. Samelson, and R.N. Germain. 1995. Zeta phosphorylation without ZAP-70 activation induced by TCR antagonists or partial agonists. *Science.* 267:515–518.
11. Kouskoff, V., S. Famiglietti, G. Lcaud, P. Lang, J.E. Rider, B.K. Kay, J.C. Cambier, and D. Nemazee. 1998. Antigens varying in affinity for the B cell receptor induce differential B lymphocyte responses. *J. Exp. Med.* 188:1453–1464.
12. Benschop, R.J., D. Melamed, D. Nemazee, and J.C. Cambier. 1999. Distinct signal thresholds for the unique antigen receptor-linked gene expression programs in mature and immature B cells. *J. Exp. Med.* 190:749–756.
13. Siskind, G.W., and B. Benacerraf. 1969. Cell selection by antigen in the immune response. In *Advanced Immunology*. F.J. Dixon and H.G. Kunkel, editors. Academic Press, New York. 1–50.
14. Hannum, L.G., D. Ni, A.M. Haberman, M.G. Weigert, and M.J. Shlomchik. 1996. A disease-related rheumatoid factor autoantibody is not tolerized in a normal mouse: implications for the origins of autoantibodies in autoimmune disease. *J. Exp. Med.* 184:1269–1278.
15. Jacob, J., R. Kassir, and G. Kelsoe. 1991. In situ studies of the primary immune response to (4-hydroxy-3-nitrophenyl)acetyl. I. The architecture and dynamics of responding cell populations. *J. Exp. Med.* 173:1165–1175.
16. Jacob, J., G. Kelsoe, K. Rajewsky, and U. Weiss. 1991. Intracloal generation of antibody mutants in germinal centres. *Nature.* 354:389–392.
17. Jacob, J., and G. Kelsoe. 1992. In situ studies of the primary immune response to (4-hydroxy-3-nitrophenyl)acetyl. II. A common clonal origin for periarteriolar lymphoid sheath-associated foci and germinal centers. *J. Exp. Med.* 176:679–687.
18. Smith, K.G., T.D. Hewitson, G.J. Nossal, and D.M. Tarlin-

- ton. 1996. The phenotype and fate of the antibody-forming cells of the splenic foci. *Eur. J. Immunol.* 26:444–448.
19. Weiss, U., and K. Rajewsky. 1990. The repertoire of somatic antibody mutants accumulating in the memory compartment after primary immunization is restricted through affinity maturation and mirrors that expressed in the secondary response. *J. Exp. Med.* 172:1681–1689.
 20. Ziegner, M., G. Steinhauser, and C. Berek. 1994. Development of antibody diversity in single germinal centers: selective expansion of high-affinity variants. *Eur. J. Immunol.* 24: 2393–2400.
 21. Bruggemann, M., A. Radbruch, and K. Rajewsky. 1982. Immunoglobulin V region variants in hybridoma cells. I. Isolation of a variant with altered idiotypic and antigen binding specificity. *EMBO J.* 1:629–634.
 22. Shlomchik, M.J., P. Watts, M.G. Weigert, and S. Litwin. 1998. Clone: a Monte-Carlo computer simulation of B cell clonal expansion, somatic mutation and antigen-driven selection. *Curr. Top. Microbiol. Immunol.* 229:173–197.
 23. Wysocki, L., T. Manser, and M.L. Gefter. 1986. Somatic evolution of variable region structures during an immune response. *Proc. Natl. Acad. Sci. USA.* 83:1847–1851.