

Differentiation of CD8 memory T cells depends on Foxo1

Rodrigo Hess Michelini,^{1,2} Andrew L. Doedens,¹ Ananda W. Goldrath,¹ and Stephen M. Hedrick^{1,2}

¹Molecular Biology Section, Division of Biological Sciences and ²Department of Cellular and Molecular Medicine, University of California, San Diego, La Jolla, CA

The forkhead O transcription factors (FOXO) integrate a range of extracellular signals, including growth factor signaling, inflammation, oxidative stress, and nutrient availability, to substantially alter the program of gene expression and modulate cell survival, cell cycle progression, and many yet to be unraveled cell type-specific responses. Naive antigen-specific CD8⁺ T cells undergo a rapid expansion and arming of effector function within days of pathogen exposure. In addition, by the peak of expansion, they form precursors to memory T cells capable of self-renewal and indefinite survival. Using lymphocytic choriomeningitis virus Armstrong to probe the response to infection, we found that *Foxo1*^{-/-} CD8⁺ T cells expand normally with no defects in effector differentiation, but continue to exhibit characteristics of effector T cells long after antigen clearance. The KLRG1^{lo} CD8⁺ T cells that are normally enriched for memory-precursor cells retain Granzyme B and CD69 expression, and fail to up-regulate TCF7, EOMES, and other memory signature genes. As a correlate, *Foxo1*^{-/-} CD8⁺ T cells were virtually unable to expand upon secondary infection. Collectively, these results demonstrate an intrinsic role for FOXO1 in establishing the post-effector memory program that is essential to forming long-lived memory cells capable of immune reactivation.

Intracellular infectious agents stimulate several thousand antigen-specific naive CD8⁺ T cells to expand up to 10,000-fold resulting in lymphocytosis and lymphadenopathy (Wirth and Harty, 2009). Within this expanded T cell population, there exist several distinct subsets that can be characterized by both function and phenotype. Cells exhibiting strong cytotoxicity to the instigating agent express high levels of perforin, granzymes, and the killer cell lectin-like receptor G1 (KLRG1). With sterilizing immunity, many of these terminally differentiated effector cells die at a high rate over a 2-wk period after the peak of the expansion. In contrast, a subset of T cells does not express KLRG1, displays a relatively reduced rate of cell death, and preferentially contributes to indelible antigen-specific immune memory (Sarkar et al., 2008; Parish and Kaech, 2009). Experiments with single-cell transfers show that these diverse populations arise from a common precursor (Stemberger et al., 2007; Gerlach et al., 2010), and this commitment may be influenced early in the process of naive T cell activation (Celli et al., 2008; Beuneu et al., 2010).

The differentiation and expansion of CD8⁺ effector T cells depends on co-stimulation, growth factors such as IL-2 (Williams et al., 2006; Bachmann et al., 2007; Obar et al., 2010; Pipkin et al., 2010), and inflammatory cytokines, especially IL-12, that promote the expression of TBX21 (Curtsinger et al., 2003; Takemoto et al., 2006; Joshi et al., 2007; Pearce and Shen, 2007). Further studies have shown that IL-2 acts, in part, through the transcriptional repressor BLIMP1 (encoded by *Prdm1*); accordingly, *Prdm1*^{-/-} T cells display a defect in effector cell differentiation and more readily form memory-precursor cells. The pro-memory transcriptional profile that includes *Bcl6*, *Tcf7*, and *Eomes* is inhibited by BLIMP1, whereas a transcription factor associated with effector T cells, TBX21, is enhanced by BLIMP1 (Kallies et al., 2009; Rutishauser et al., 2009; Shin et al., 2009; Ji et al., 2011; Yang et al., 2011).

© 2013 Hess Michelini et al. This article is distributed under the terms of an Attribution-Noncommercial-Share Alike-No Mirror Sites license for the first six months after the publication date (see <http://www.rupress.org/terms>). After six months it is available under a Creative Commons License (Attribution-Noncommercial-Share Alike 3.0 Unported license, as described at <http://creativecommons.org/licenses/by-nc-sa/3.0/>).

CORRESPONDENCE

S. M. Hedrick:
shedrick@ucsd.edu

Abbreviations used: KLRG1, killer cell lectin-like receptor G1; LCMV, lymphocytic choriomeningitis virus.

Studies have shown that the AKT signaling pathway promotes effector cell differentiation at the expense of memory cell precursors (Hand et al., 2010; Kim et al., 2012). In addition, the mammalian target of rapamycin, a downstream target of AKT, is a major regulator of memory CD8⁺ T cell differentiation (Araki et al., 2009; Pearce et al., 2009). Thus treatment with rapamycin or metformin enhanced both the quantity and quality of memory CD8⁺ T cells. Because inhibition of the FOXO1 transcription factor is a major conduit of AKT-mediated signaling, we set out to determine whether FOXO1 broadly affects the contingency of effector versus memory-precursor differentiation, and to what extent FOXO1 determines the program of memory T cell gene expression. Here, we show that the loss of *Foxo1* has little effect on the expansion and survival of antigen-stimulated CD8⁺ T cells, but causes them to maintain an activated effector phenotype. These persisting *Foxo1*-null T cells were unable to expand upon reactivation, and this phenotype correlated with a extensive pattern of gene expression that favors the formation of effector T cells. We conclude that FOXO1 regulates the fate of effector versus memory-precursor T cells, and this has implications for the manner with which the physiological state of the organism impacts the outcome of an immune response.

RESULTS

Normal expansion of *Foxo1*^{-/-} CD8⁺ T cells after viral infection

To study the role of FOXO1 in CD8⁺ T cells responding to a viral infection in vivo, we took advantage of mice in which exon 2 of *Foxo1* is flanked by *loxP* sites (*Foxo1*^{fl/fl} mice; Paik et al., 2007). Because crossing *Foxo1*^{fl/fl} mice with mice carrying the Tg(Cd4-cre)1Cwi transgene (Cd4Cre mice) results in autoimmunity caused by a defect on regulatory T cells (Kerdiles et al., 2010), we produced *Foxo1*^{fl/fl} mice with the Cre recombinase under the control of the human Granzyme B promoter (GZMBCre). Recombination was thus restricted to activated CD8⁺ T cells (Jacob and Baltimore, 1999; Rutishauser et al., 2009). These mice were further crossed to include the P14 T cell antigen receptor transgenes (P14) endowing T cells with specificity for the immunodominant epitope of lymphocytic choriomeningitis virus (LCMV) gp33 in association with H2D^b. Equal numbers of spleen cells from *Foxo1*^{fl/fl}GZMBCre⁺P14⁺ (*Foxo1*^{-/-} P14) and *Foxo1*^{fl/fl}GZMBCre⁻P14⁺ (WT P14) mice were co-transferred into naive host mice, and the T cells were followed by expression of CD45.2 or CD45.1/2, respectively. Recipient mice were then infected with LCMV Armstrong (LCMV-Arm; Oldstone, 2002). The analysis of splenic T cells revealed an equivalent expansion of *Foxo1*^{-/-} and WT P14 T cells upon viral infection with little or no difference in the rate of contraction (Fig. 1 A). Although there appears to be a slight delay in the contraction of *Foxo1*^{-/-} T cells in the experiment presented, this was not consistently seen in three other independent experiments. In addition, similar percentages of Ki-67⁺ cells in WT and *Foxo1*^{-/-} P14 cells were found at the peak of the response (7 d after infection) with virtually no Ki-67⁺ cells present at days 12 and 19 (Fig. 1 B, top panels and bottom

graph). Since FOXO1 is important for the expression of the homing molecules, L-selectin (CD62L) and CCR7 (Fabre et al., 2008; Kerdiles et al., 2009), we examined CD62L expression in the spleen and the distribution of WT and *Foxo1*^{-/-} P14 cells after viral infection in lymph nodes (LN), lungs, and liver. As expected, expression of CD62L was low on *Foxo1*^{-/-} P14 cells compared with WT P14 cells (Fig. 1 C). However, *Foxo1*^{-/-} T cells were at least as abundant as WT P14 T cells in all the organs analyzed, excluding a defect in the homing capacity of *Foxo1*^{-/-} P14 cells due to low CD62L expression (Fig. 1 D). Thus, loss of FOXO1 had no effect on the overall proliferation of virus-specific CD8⁺ T cells in response to LCMV infection.

FOXO transcription factors are subject to complex post-translational regulation that includes nuclear egress and cytoplasmic localization followed by 14-3-3-mediated degradation. As such, we sought to determine how the amounts of FOXO1 might change during the course of CD8⁺ T cell expansion and contraction. The results showed there was heterogeneous FOXO1 expression that was inversely correlated with KLRG1 expression (Fig. 1 E, WT). We also note that FOXO1 expression was not detected in the GZMBCre⁺ T cells. These results are consistent with the possibility that FOXO1 is active in the precursors to memory cells, and plays less of a role in KLRG1^{hi} cells. To address this issue, we focused our analysis on memory CD8⁺ T cell differentiation with or without the deletion of *Foxo1*.

KLRG1^{lo} CD8⁺ memory-precursor T cells display an enhanced effector phenotype

After the peak expansion of CD8⁺ T cells in response to virus infection, effector cells selectively die, but the surviving effector cells and memory-precursor cells become quiescent, and those within the secondary lymphoid organs eventually extinguish the expression of cytotoxic effector molecules such as GZMB (Kaech et al., 2002). To examine how the deletion of *Foxo1* affected the resolution of the response, we examined CD8⁺ T cells for the expression of molecules that play a role in effector function. Although the percentage of GZMB⁺ cells among WT and *Foxo1*^{-/-} P14 cells was similar at the peak of the response (Fig. 2 A, top left), the mean fluorescence intensity (MFI) in *Foxo1*^{-/-} P14 cells was higher (Fig. 2 A, top right). In addition, WT P14 cells rapidly lost GZMB during the contraction phase (d12-19), whereas *Foxo1*^{-/-} P14 cells retained GZMB in reduced, though substantial amounts. In fact, the KLRG1^{lo} cells increased GZMB expression at d19 against a constant expression within the KLRG1^{hi} population. Similarly, the KLRG1^{lo} *Foxo1*^{-/-} T cells regained expression of CD69 during the contraction phase (Fig. 2 A, bottom), and comparable results for the expression of GZMB and CD69 were found for liver T cells (not depicted). These data suggest that in the absence of FOXO1, CD8⁺ T cells fail in their transition from effector to memory cells, showing a sustained effector phenotype. How the CD8⁺ T cells reexpress effector molecules in the absence of antigen is unclear. In accord with the enhanced and extended effector function, there was a greater number of *Foxo1*^{-/-} multicytokine-producing (TNF and IFN- γ) effector T cells compared with WT (Fig. 2 B).




Figure 1. Kinetics of CD8⁺ T cell expansion in response to LCMV-Arm infection. A 1:1 mix of 10⁴ CD45.1^{+/2+} WT (*Foxo1*^{fl/fl}GZM-B^{Cre}-) and CD45.2⁺ *Foxo1*^{-/-} (*Foxo1*^{fl/fl}GZM-B^{Cre}+) P14 cells were transferred to CD45.1⁺ WT host mice. Mice were then infected with LCMV-Arm, and lymphoid and nonlymphoid organs were harvested and analyzed at day 7, 12, and 19 after infection. (A) WT and *Foxo1*^{-/-} P14 cells were identified in the spleen by their expression of CD45.1, CD45.2, CD8, and V α 2 and total numbers of P14 cells were determined. DPI, days post infection. (B) Ki67 staining of spleen cells from WT and *Foxo1*^{-/-} P14 cells at different time points. (C) Expression of KLRG1 and CD62L on WT and *Foxo1*^{-/-} P14 cells isolated from the spleen. (D) Total numbers of WT and *Foxo1*^{-/-} P14 cells in the indicated organs. (E) Expression of KLRG1 and FOXO1 on WT and *Foxo1*^{-/-} P14 cells isolated from the spleen. Data representative of 1 out of 3 independent experiments each, with $n = 3,4$.

To ensure that *Foxo1*^{-/-} T cells were functional, we infected WT (*Foxo1*^{fl/fl}GZM^BCre⁻) and *Foxo1*^{-/-} (*Foxo1*^{fl/fl}GZM^BCre⁺) mice and found no detectable LCMV virus by day 7 (unpublished data).

At the peak of the expansion phase, T cell diversity is evident based on the differential expression of several diverse cell surface receptors and transcription factors. For example, the KLRG1^{hi}CD127^{lo} population is enriched for committed effector cells, whereas the KLRG1^{lo}CD127^{hi} population is enriched for the precursors of memory T cells (Joshi et al., 2007). A complication in the present analysis is that FOXO1 directly

controls the expression of CD127 at least in naive and acutely activated T cells (Kerdiles et al., 2009), and as such may not accurately reflect the state of CD8⁺ T cell differentiation. This is depicted in Fig. 3A, at 7 and 12 d, where an emerging CD127⁺ population is visible in the WT but not the *Foxo1*^{-/-}. However, we note that by 19 d, CD127 expression has been reestablished on a proportion of the KLRG1^{lo} cells, and this suggests that at least a subset of activated T cells can express CD127 in the absence of FOXO1.

The T-box transcription factors EOMES (eomesodermin) and TBX21 (T-bet) are required for the formation and function




Figure 2. Phenotype of virus-specific CD8⁺ T cells.

A 1:1 mix of 10⁴ CD45.1⁺/2⁺ WT (*Foxo1*^{fl/fl}GZMBCre⁻) and CD45.2⁺ *Foxo1*^{-/-} (*Foxo1*^{fl/fl}GZMBCre⁺) P14 cells were transferred to CD45.1⁺ WT host mice. Mice were then infected with LCMV-Arm, and spleen cells were harvested and analyzed at different time points. (A) The expression of KLRG1, GZMB, and CD69 was determined in WT and *Foxo1*^{-/-} P14 cells at different time points (left), and the amount of GZMB (MFI) and percentage of CD69 are depicted (right). Data are representative of 3 independent experiments, with *n* = 3–4. (B) At day 7 after infection, spleen cells were stimulated with PMA and ionomycin and the frequency of IFN-γ- and TNF-producing cells was determined. Data are representative of 2 independent experiments each, with *n* = 4.

of effector and memory CD8⁺ T cells (Kaech and Cui, 2012). TBX21 is required for the formation of KLRG1^{hi} effector cells, and its overexpression skews the population toward terminal effector cell differentiation. In contrast, EOMES is required for T cells to survive in the long term as memory cells expressing IL2Rβ (CD122; Intlekofer et al., 2005; Banerjee et al., 2010). An examination of co-transferred WT and *Foxo1*^{-/-} P14 cells showed that within the WT T cell population there was a remaining KLRG1^{lo}EOMES^{hi} population at 19 d after infection that was largely missing in the *Foxo1*^{-/-} P14 cells (Fig. 3 B). This correlates with the absence of the high-mobility group protein, TCF7, a Wnt-activated transcription factor shown to be important for *Eomes* expression and maintenance of CD8⁺ memory T cells (Zhou et al., 2010; Fig. 3 C). Thus, there was a substantial deficit in the expression of two transcription factors shown to be required for the maintenance and function of CD8⁺ memory T cells.

Collectively, these results demonstrate an important role for FOXO1 in regulating essential characteristics associated with the lineage of precursor cells destined to become self-renewing,

long-lived memory T cells. A recent study showed that IL-12 promotes *Tbx21* expression by inactivating FOXO1, and *Tbx21* expression can be inhibited indirectly by FOXO1 overexpression (Rao et al., 2012). Furthermore, FOXO1 appears to directly target *Eomes*, and thus constitutes a second pathway that converges on *Eomes* expression to promote a memory cell phenotype (Rao et al., 2012). In line with these results, *Foxo1*^{-/-} P14 cells displayed higher TBX21 expression compared with WT P14 cells (Fig. 3 D), and this correlated with lower levels of EOMES (Fig. 3 B), supporting a critical role for FOXO1 in regulating both *Tbx21* and *Eomes* expression.

Finally, CXCR3 is up-regulated on CTLs after activation and is thought to facilitate colocalization of T cells with viral antigens in the spleen. Whereas KLRG1^{hi} P14 cells were shown to exhibit a heterogeneous expression pattern, KLRG1^{lo} T cells remained uniformly CXCR3^{hi} throughout a viral response (Hu et al., 2011). This was seen for both WT and *Foxo1*^{-/-} P14 cells (Fig. 3 E), suggesting that *Foxo1*^{-/-} CD8⁺ T cells have equal access to antigen, and that this aspect of CTL differentiation appears to be unaffected by *Foxo1* deletion.




Figure 3. Enhanced effector phenotype in the absence of FOXO1. A 1:1 mix of 1×10^4 CD45.1^{+/2+} WT (*Foxo1^{fl/fl}GZMBCre⁻*) and CD45.2⁺ *Foxo1*^{-/-} (*Foxo1^{fl/fl}GZMBCre⁺*) P14 cells were transferred to CD45.1⁺ WT host mice. Mice were then infected with LCMV-Arm, and spleen cells were harvested and analyzed at different time points. (A–E) WT, open bars; *Foxo1*^{-/-}, filled bars. (A) The expression of KLRG1 and CD127 was determined in WT and *Foxo1*^{-/-} P14 cells, and the percentage of CD127⁺ cells is depicted (bottom). (B) The expression of KLRG1 and EOMES was determined in WT and *Foxo1*^{-/-} P14 cells, and the percentage of EOMES⁺ cells is depicted (bottom). (C and D) WT and *Foxo1*^{-/-} P14 cells were analyzed for their expression of TCF7 (C) and TBX21 (D), and the amount of both are presented as mean fluorescence intensity (MFI, right). (E) The expression of KLRG1 and CXCR3 was determined in WT and *Foxo1*^{-/-} P14 cells (left), and the percentage of CXCR3⁺ cells depicted (right). Data are representative of 3 independent experiments each, with $n = 3-4$.

Foxo1^{-/-} CD8⁺ T cells fail to up-regulate memory-associated signature genes

To obtain an unbiased analysis of genes differentially expressed in antigen-specific *Foxo1*^{-/-} CD8⁺ T cells responding to infection, we performed microarray analysis on KLRG1^{lo} and KLRG1^{hi} FACS-sorted, congenically marked WT and *Foxo1*^{-/-} P14 cells 8 d after infection with LCMV-Arm (Fig. 4). For this experiment, we performed gene deletion in *Rosa26^{Cre-ERT2} Foxo1^{fl/fl}* (*Foxo1*^{-/-}) P14 mice just before adoptive transfer (Kerdiles et al., 2009). A virtually complete deletion of *Foxo1* was obtained before adoptive transfer and maintained throughout the duration of the experiment (unpublished data). We used

a replicate mixed transfer experimental design to sort *Foxo1*^{fl/fl} (WT) and *Foxo1*^{-/-} P14 cells of the same TCR specificity from the same infected animals, thus controlling for cytokine environment and antigen load on CTL gene expression (Fig. 4 A). We compared the gene expression profile between WT versus *Foxo1*^{-/-} P14 CD8⁺ T cells in KLRG1^{lo} (Fig. 4 B) and KLRG1^{hi} (Fig. 4 C) sorted cells. The data showed that FOXO1 regulates gene expression in both populations. We then focused our analysis on sorted KLRG1^{lo} T cells known to seed the memory population and display abundant differences in both effector molecules and key transcription factors when compared with KLRG1^{hi} T cells (Figs. 2 and 3, respectively;




Figure 4. Deletion of *Foxo1* prevents the development of a memory program of gene expression. A 1:1 mix of 10⁴ WT (*Foxo1*^{fl/fl} *Rosa26*) and *Foxo1*^{-/-} (*Foxo1*^{fl/fl} *Rosa26*^{Cre-ERT2}) P14 cells were prepared from tamoxifen-treated mice and transferred to CD45.1⁺ WT host mice. Microarray gene-expression analysis was performed by sorting WT and *Foxo1*^{-/-} KLRG1^{lo} P14 cells 8 d after infection with LCMV-Arm. (A) Experimental scheme for mixed transfer and FACS of WT and *Foxo1*^{-/-} P14 T cells on day 8 of LCMV-Arm infection. The four double-sorted populations were analyzed on the same

Kaech and Cui, 2012). We identified genes passing a twofold up or down cut-off in KLRG1^{lo} cells, identifying 27 genes down and 59 genes up in WT relative to *Foxo1*^{-/-} P14 CD8⁺ T cells (Fig. 4, B and D). As expected, and as a positive control for our analysis, we observed decreased abundance of mRNAs of known FOXO1 targets in *Foxo1*^{-/-} P14 CD8⁺ T cells: *Sell* (3.9 fold); *Ctla4* (3.8 fold); *Il7r* (10.1 fold); and *Foxo1* itself (2.5 fold; Fig. 4 D). We grouped the twofold list of regulated genes into naive/memory, effector, or no pattern based on their expression during the CD8⁺ T cell response to infection as reported by the Immgen project (Best et al., 2013). Interestingly, *Foxo1*^{-/-} P14 CD8⁺ T cells were enriched for genes associated with effector T cells, whereas WT P14 CD8⁺ T cells genes were enriched for genes associated with T cell naive/memory (Fig. 4 D, pie charts).

Although 8 d after infection has been found to be the peak of cytotoxic T cell expansion, virus has been cleared by this point, and subsets of CTL can already be identified as terminally differentiated effector cells or memory cell precursors. Memory precursors are associated with expression of *Id3*, *Eomes*, and *Tcf7*, whereas terminally differentiated effectors are associated with expression of *Prdm1* (BLIMP1), *Id2*, and elevated *Tbx21* (Kaech and Cui, 2012). As shown, these signature transcription factors are differentially regulated in KLRG1^{lo} *Foxo1*^{-/-} versus WT P14 cells (Fig. 4 E, left); furthermore, these results are consistent with aforementioned results (Fig. 3, B–D) showing that TCF7 and EOMES were markedly reduced in *Foxo1*^{-/-} T cells, whereas *Tbx21* was increased. In addition, CTLs exert their effector function through the granule exocytosis pathway, FAS, and cytokine production. We found elevated granzymes, perforin, *Fas*, and *Ifng* in KLRG1^{lo} *Foxo1*^{-/-} P14 cells. We further analyzed KLRG1^{lo} cells for activation and trafficking molecules (Fig. 4 E, right). Altogether, these data show a bias for *Foxo1*^{-/-} CD8⁺ T cells toward terminal differentiation, increased effector function, and altered trafficking/activation (Fig. 4 E, middle and right).

From our sorted cells, we made an expression list of genes up or down in WT KLRG1^{lo} versus WT KLRG1^{hi}. This established a list of genes expressed at higher levels in terminal-effector cells relative to those T cells with increased memory potential. When we overlaid the genes up or down 1.5-fold from this list into a volcano plot of WT versus *Foxo1*^{-/-} KLRG1^{lo} CD8⁺ T cells, there was an ~90% concordance of transcripts; genes associated with terminal effector cells were

increased in *Foxo1*^{-/-} T cells, whereas genes associated with memory-precursor cells were increased in WT (Fig. 4 F). Specifically, the results show that *Foxo1*^{-/-} KLRG1^{lo} cells are enriched for terminally differentiated transcripts (87% of KLRG1^{hi}-biased transcripts are higher in KLRG1^{lo} *Foxo1*^{-/-} versus KLRG1^{lo} WT), whereas WT KLRG1^{lo} cells are enriched for memory transcripts (88% of the KLRG1^{lo}-biased transcripts are up in WT KLRG1^{lo} versus *Foxo1*^{-/-} KLRG1^{lo}). A trivial explanation for this analysis would be that there was a bias for KLRG1^{hi} cell differentiation caused by the absence of FOXO1, and the microarray data simply reflects this while not revealing a direct role for *Foxo1* in driving memory cell differentiation. However, at the peak of the response there are approximately equivalent proportions of KLRG1^{hi} and KLRG1^{lo} cells in WT and *Foxo1*^{-/-} T cells, and yet TCF7, EOMES, TBX21, and GZMB on KLRG1^{lo} cells are differentially expressed within these populations throughout infection (Figs. 2, 3). We infer that the effect is not simply a population bias but an intrinsic FOXO1-dependent effect on key memory and effector molecules. These results suggest a failure of *Foxo1*^{-/-} CD8⁺ T cells at the peak of the response to differentiate into memory-precursor cells.

The fact that so many genes associated with the memory state are of decreased abundance in *Foxo1*^{-/-} T cells suggests two nonmutually exclusive possibilities. One, FOXO1 is essential to terminate the effector state allowing genes required for differentiated memory cells to be expressed under the control of other factors. Two, FOXO1 is essential for driving the memory program through direct induction of known targets *Sell*, *Il7r*, and of master transcription factors of memory such as *Tcf7* and *Eomes* (Rao et al., 2010; Zhou et al., 2010). *Tcf7* is expressed in both CD4⁺ and CD8⁺ T cells from early thymic developmental stages and in peripheral cells, it is notably absent in effector cells and then once again expressed in high amounts in memory T cells (Yu et al., 2010; Best et al., 2013).

In support of a direct role for FOXO1 in memory cell formation, we have determined, using a chromatin immunoprecipitation sequencing (ChIP-Seq) approach, that FOXO1 binds directly to multiple sites close to, and within the *Tcf7* gene in naive CD4⁺ T cells, and these binding sites were also found in two other T cell-specific ChIP-Seq datasets (Fig. 4 G). Five of the peaks were tentatively labeled as E (intergenic enhancer), P (promoter), I1 (intronic enhancer-1), I2 (intronic enhancer-2), and I3 (intronic enhancer-3). For each of these elements, we list the nucleotide sequences corresponding to each ChIP-Seq peak

microarray chip. (B) KLRG1^{lo} WT versus *Foxo1*^{-/-} reveals 59 genes up and 27 genes down with a twofold cutoff. (C) KLRG1^{hi} WT versus *Foxo1*^{-/-} reveals 47 genes up and 45 genes down with a twofold cutoff. (D) Identification of genes KLRG1^{lo} WT / *Foxo1*^{-/-} with a twofold cutoff. Pie charts indicate portion of genes with an expression pattern tied to naive/memory (blue), effector (red), or no pattern (gray). Asterisk indicates expression was verified by flow cytometry. (E) Classification of a subset of the genes in (D) by function. (F) Genes differentially regulated in WT KLRG1^{lo} versus WT KLRG1^{hi}, were overlaid with genes from KLRG1^{lo} WT versus KLRG1^{lo} *Foxo1*^{-/-} CD8⁺ T cells, showing nearly a 90% concordance of transcripts. (G) Naive CD4⁺ ChIP-Seq data showing several FOXO1 binding sites proximal to and within *Tcf7*. Data were analyzed from T reg cells and two different datasets analyzing naive CD4 T cells. Peaks are labeled as putative: intergenic enhancer (E), promoter (P), and intronic enhancers 1, 2, and 3 (I1, I2, I3). Peak sequences are listed in Table 1. (H) Transcription factors known to play a role in the formation of memory cells, summarizing data presented in D with the addition of other transcription factors differentially regulated in WT vs. *Foxo1*^{-/-} P14 cells. Data are from 1 experiment with $n = 3$.

(Table I). As shown, each has one or more tandem consensus forkhead binding sequences at or surrounding the peak, and this is consistent with specific FOXO1 binding. In addition, we show the H3K4me3 histone marks at the transcription start site indicative of an open locus (Pekowska et al., 2011). This apparently direct regulation of *Tcf7* suggests that memory cell differentiation after TCR and cytokine stimulation requires FOXO1 nuclear localization.

Finally, for clarity, we summarized the data focusing on transcription factors differentially regulated in WT versus *Foxo1*^{-/-} P14 T cells (Fig. 4 H). This list includes other transcription factor genes known to play a role in the formation of memory, such as the *Tcf7*-related *Lef1* (Zhou and Xue, 2012). It also may point to a role for FOXO1-regulated transcription factors and DNA binding proteins as of yet not recognized as being involved in CD8⁺ T cell memory formation.

Collectively, these analyses suggest a basic and essential role for FOXO1 in programming fundamental aspects of the memory state in CD8⁺ T cells after response to infection in vivo. We therefore tested whether *Foxo1*^{-/-} T cells can expand in response to a secondary antigenic challenge as a measure of the requirement for *Foxo1* in an immune recall response.

Foxo1^{-/-} CD8⁺ T cells fail to make a memory response

The data presented demonstrate that, in the absence of FOXO1, virus-specific CD8⁺ T cells display a prolonged effector phenotype and fail to up-regulate a multitude of key memory-associated genes. To better understand the role of FOXO1 in CD8⁺ T cell differentiation after viral infection, we characterized *Foxo1*^{-/-} P14 spleen cells from LCMV-Arm infected mice at later time points (35 d after infection). As seen for early time points (Fig. 1 A), the numbers of P14 cells were comparable between WT and *Foxo1*^{-/-} P14 cells (Fig. 5 A). Even at this later time, *Foxo1*^{-/-} P14 cells exhibit an enhanced effector-like phenotype (Fig. 5 B) characterized by low expression of the memory-associated marker CD127, and high expression of CD69 and GZMB. As shown for early time points (Fig. 2 A; and Fig. 3, A and B), this phenotype was only

Table 1. Nucleotide sequences found at FOXO1-binding peaks within the *Tcf7* gene

Element ^a	Peak sequences
E	5'-TATCTGTTTTGCTC-3'
P	5'-GGAGTAAACAGACCC-3'
I1	5'-ACTGTTGTTTCCTGC-3' 5'-TGTACAAACAAGGCT-3' 5'-GGAGGAAACAGGTGT-3'
I2	5'-CAGGGTGTTTGTAGT-3' 5'-TCTAAAAACATCCTG-3' 5'-AAGGAAAACACAAGC-3'
I3	5'-GACTGTGTTTATTTT-3' 5'-CTCTGAAACAGAGAC-3'

Sequences in bold correspond to a forkhead consensus site.

^aSee Fig. 4 G.

observed in KLRG1^{lo} memory-like cells (Fig. 5, B and C). This KLRG1^{lo} population also expressed high amounts of the high-affinity IL-2 receptor CD25 and low levels of the IL-2 receptor β -chain CD122 (Fig. 5 C), consistent with the microarray data (Fig. 4 E, right). Early in infection, high CD25 expression was shown to promote terminal-effector CD8⁺ T cell differentiation in vivo (Kalia et al., 2010), while high CD122 expression is associated with CD8⁺ memory T cells (Intlekofer et al., 2005). This supports the concept that FOXO1 is essential for the transition from effector to long-lived memory CD8⁺ T cells. In addition, the expression of memory-related transcription factors EOMES and TCF7 was still low, whereas the expression of TBX21 was higher in KLRG1^{lo} *Foxo1*^{-/-} P14 cells compared with WT at this late time-point (Fig. 5 D). These phenotypic changes are consistent with an extended effector-like phenotype in *Foxo1*^{-/-} T cells (Fig. 5 B).

Finally, we tested the ability of these cells to respond to a second challenge with LCMV-Arm. To do this, WT and *Foxo1*^{-/-} P14 cells were sorted from mixed adoptively transferred mice at 35 d after infection and transferred as a 1:1 mixture into naive recipients. These mice were challenged with LCMV-Arm and analyzed for T cell expansion 5 d post-infection (Fig. 5 E). The results showed that in the absence of FOXO1, CD8⁺ T cells failed to accumulate, and were at a 45-fold disadvantage with respect to the WT CD8⁺ T cells. Of note, the phenotype of the cells post-challenge (Fig. 5 E) was similar to the one found at the peak of the response during the primary infection (Figs. 2 and 3), with higher expression of GZMB in *Foxo1*^{-/-} P14 cells compared with WT P14 cells. Collectively, the results of the experiments described demonstrate an intrinsic role for FOXO1 in regulating the transition from effector cells to long-lived memory cells capable of reactivation.

DISCUSSION

Infection by an intracellular pathogen almost invariably results in the expansion of antigen-specific CD8⁺ T cells that participate in killing the pathogen-infected cells. These T cells exhibit characteristics of terminal differentiation in that they have a finite life span and they maintain a highly differentiated program of gene expression associated with cytotoxicity and inflammation. Although competing models of CTL memory exist, available evidence favors the concept that memory-precursor cells differentiate from a subset of this effector population, ultimately becoming a cell-type perhaps like no other in the immune system. These memory T cells are self-renewing, with a population turnover of 2–3 mo, and their progeny can be shown to persist for the life of the individual (Parretta et al., 2008; Choo et al., 2010). Yet, they are poised to respond to a reinfection by rapid growth, progression through the cell cycle, and expression of a cytotoxic, pro-inflammatory program of gene expression (Tanchot et al., 1997; Murali-Krishna et al., 1999; Zhang and Bevan, 2011).

The control of the effector versus memory-precursor lineage decision has been illuminated over the past decade, including a role for stimulatory and inflammatory cytokines as well as a description of several transcription factors necessary for the distinct programs of gene expression (Rutishauser and

Kaech, 2010; Best et al., 2013). In this study, we describe experiments that demonstrate a high order control over this fate decision mediated by the forkhead transcription factor, FOXO1. With the deletion of *Foxo1*, CD8⁺ T cells fail to acquire the characteristics of memory-precursor cells 7 d post-infection, they retain many aspects of effector T cells for an extended period, and they do not exhibit the vast expansion characteristic of a recall immune response. Given its extensive range of posttranslational modifications (Hedrick et al., 2012), we infer that the lineage decision to acquire characteristics of effector vs. memory-precursor cells would appear to center, to an extent, on the integration of extrinsic signals that are manifest in the cellular localization and target specificity of FOXO1.

These results are in concert with studies showing that activation of AKT, a major pathway to FOXO inhibition, is key to the differentiation of effector cells at the expense of memory-precursor cells (Hand et al., 2010; Kim et al., 2012). In addition, the IL-12-mediated inactivation of FOXO1 was shown to promote TBX21-expressing effector T cells, and the rapamycin-mediated inhibition of TBX21 expression was FOXO1 dependent. Also, this same study established a role for FOXO1 in switching T cells from an effector to a memory-like phenotype through an increase in EOMES expression (Rao et al., 2012). Here, we establish that FOXO1, in the context of rapidly expanding T cells, directly and indirectly refines a program of gene expression to produce the phenotypic characteristics of memory cells. Of particular importance




Figure 5. *Foxo1*^{-/-} CD8⁺ T cells fail to expand upon secondary viral challenge. (A–E) WT, open squares; *Foxo1*^{-/-} filled squares. A 1:1 mix of 10⁴ CD45.1^{+/2+} WT (*Foxo1*^{fl/fl}GZMBCre^{-/-}) and CD45.2^{+/2+} *Foxo1*^{-/-} (*Foxo1*^{fl/fl}GZMBCre^{+/+}) P14 cells were prepared and transferred to CD45.1⁺ WT host mice. Mice were then infected with LCMV-Arm, and spleen cells were analyzed 35 d after infection. (A) Number of cells recovered per spleen at day 35. (B) WT and *Foxo1*^{-/-} P14 cells were analyzed by their expression of KLRG1 and CD127, CD69, and GZMB (top), and the MFI of these molecules was quantified on the KLRG1^{lo} population (bottom). (C) Expression of CD25 and CD122 on KLRG1^{lo} P14 cells. (D) WT and *Foxo1*^{-/-} P14 cells were analyzed for expression of transcription factors important for CD8⁺ memory and effector T cells, and the amount of protein expressed on the KLRG1^{lo} population graphed as MFI. (E) P14 (CD8⁺Vα2⁺) WT (CD45.1^{+/2+}) and *Foxo1*^{-/-} (CD45.1⁺) cells from hosts 35 d after LCMV-Arm infection were sorted and transferred at a 1:1 ratio into naive WT mice. Mice were then infected with LCMV-Arm and the expansion of WT and *Foxo1*^{-/-} P14 cells determined at d 5 after infection (top right). The phenotype of P14 T cells 5 d after secondary challenge (bottom). Data are representative of 2 independent experiments for a total *n* = 11.

may be a direct role for FOXO1 in controlling the expression of *Tcf7*, encoding a Wnt, β -catenin-regulated transcription factor. This might be viewed as contrary to studies showing that FOXO factors compete with β -catenin for TCF interactions, and thus suppress the expression of TCF target genes; however, these studies examined FOXO3 and FOXO4. We surmise that this effect may not apply to FOXO1, but rather, FOXO1 may transcriptionally promote the expression of *Tcf7* (Almeida et al., 2007; Hoogeboom et al., 2008). In this regard, the present results closely parallel those analyzing *Tcf7* mutant T cells in that such T cells maintain expression of GZMB, and exhibit reduced expression of EOMES, CD62L, CCR7, and IL2 β (Zhou et al., 2010). Our conclusion is that FOXO1 is required for the expression of *Tcf7* as part of an extended program of gene expression that gives rise to memory T cells.

FOXO1 has been found to regulate pluripotency and self-renewal in several contexts. In embryonic stem cells (ES cells) in mouse and human beings, it does so by direct control of *Oct4* and *Sox2*—encoding two of the factors necessary to maintain pluripotency in ES cells or induce pluripotency in differentiated cell types (Zhang et al., 2011). It is necessary for the maintenance of tumor stem cells in acute myelogenous leukemia (AML) cells, which explains the paradoxical finding that AKT activation or the inactivation of multiple *Foxo* genes can inhibit tumorigenicity in many examples of AML (Sykes et al., 2011). In addition, FOXO factors are essential for the maintenance of hematopoietic stem cells, although largely through the detoxification of reactive oxygen species (Tothova et al., 2007). Although we might be tempted to draw parallels between the self-renewal capacity of stem cells and memory T cells, we note that FOXO transcription factors have also been shown to control acne (Melnik, 2011), bone homeostasis (Almeida, 2011), liver metabolism (Kousteni, 2012), muscle mass (Goodman et al., 2011), and the trade-off between aging and cancer (Kloet and Burgering, 2011). In fact, there may be few physiological processes that are not guided, at least in part, by FOXO transcription factors. Perhaps what all these processes have in common is a requirement for integrating characteristics defining the physiological state of the organism to effect an adaptively advantageous program of gene expression different for each and every tissue. A corollary to this thesis is that gene expression is not determined by any one transcription factor; rather, a transcription factor exerts activity in a highly context-dependent manner (Barolo and Posakony, 2002).

These results imply that AKT-mediated inhibition of FOXO1 ensures the memory program is not prematurely engaged, as early in the response, the cytokine or TCR-driven cytoplasmic sequestration of FOXO1 may promote high TBX21 and an effector phenotype so long as antigen is present. How this pathway plays into T cell exhaustion in chronic infection or continued formation of memory T cells in viral latency will be a topic of future studies. An implication is that the magnitude and effectiveness of a recall immune response may depend on the physiological state of the individual, where high levels of growth factors or superabundant nutrients may well blunt the formation or function of memory T cells. In addition, the role of memory

T cells in persistent or latent viral infections is only beginning to be understood (Youngblood et al., 2012), and the course of such an infection could be readily affected by the activity of FOXO1 in antigen-specific T cells. The delicate balance between latency and recrudescence may be controlled in part through inactivation of FOXO caused by physiological stress.

MATERIALS AND METHODS

Mice and tamoxifen treatment. C57BL/6 CD45.1⁺ mice were maintained in pathogen-free conditions. *Rosa26*^{Cre-ERT2} mice were provided by T. Ludwig (The Ohio State University, Columbus, OH; Guo et al., 2007). *Foxo1*^{f/f} and Tg(GZMB-cre)1Jcb/J (GZMBCre) mice have been previously described (Jacob and Baltimore, 1999; Paik et al., 2007). *Foxo1*^{f/f} mice (backcrossed to C57BL/6, *n* > 12) were crossed to GZMBCre or *Rosa26*^{Cre-ERT2} mice. In addition, mice were crossed with Tg(TcrLCMV)327Sdz (P14) mice to generate *Foxo1*^{f/f} GZMBCre⁺P14 or *Foxo1*^{f/f} *Rosa26*^{Cre-ERT2} P14 CD45.2⁺ mice. CD45.1⁺/2⁺ WT P14 mice were produced by a further cross with CD45.1⁺ C57BL/6 mice. *Rosa26*^{Cre-ERT2}-mediated deletion of floxed alleles was induced by intraperitoneal injection of 1 mg of tamoxifen (Sigma-Aldrich) emulsified in 200 ml of sunflower seed oil (Sigma-Aldrich) every day for 5 d, followed by 5 d of rest (*ER*^{Cre+}, *Foxo1*^{-/-}; *ER*^{Cre-}, WT). All procedures were approved by the Animal Care and Use Committee of the University of California, San Diego.

Adoptive transfer and viral infection. Spleen cells were harvested from mice as described in the previous section. P14 cells were identified as CD8⁺V α 2⁺ and numbers were adjusted to obtain a 1:1 mix of 10⁴WT:*Foxo1*^{-/-} cells. P14 cells were transferred intravenously to 8–10-wk-old CD45.1⁺ WT host mice. Mice were then infected with 2 \times 10⁵ pfu LCMV-Armstrong i.p. Virus was grown, identified, and quantified as previously described (de la Torre and Oldstone, 1992).

Flow cytometry. Cell suspensions isolated from the indicated organs were incubated for 15 min at 4°C in PBS containing 1% FCS (Omega), 2 mM EDTA, 0.01% NaN₃, and the indicated fluorochrome-conjugated antibodies. All intracellular staining was done with FoxP3 Fix/Perm (eBioscience). Antibodies against FOXO1 and TCF7 (rabbit polyclonal) were purchased from Cell Signaling Technology. Alexa Fluor 647 anti-rabbit secondary antibody and PE-conjugated anti-human Granzyme B (GZMB) antibodies were purchased from Invitrogen. The rest of the antibodies used were purchased from eBioscience or BD. Data were collected on a Fortessa (BD) and analyzed with FlowJo software (Tree Star).

Cell sorting and microarray. Splenocytes were harvested and the CD8⁺ fraction enriched by negative selection using biotinylated antibodies against CD4, B220, CD11c, and DX5 (all from eBioscience), followed by streptavidin microbeads using MACS system in a 4°C cold room (Miltenyi Biotec). The negative fraction was then stained with antibodies against CD45.1, CD45.2, KLRG1, CD8, and V α 2. KLRG1^{lo} donor WT (CD45.1⁺/2⁺) and *Foxo1*^{-/-} (CD45.2⁺) P14 (CD8⁺V α 2⁺) splenocytes were sorted from host mice on day 7 after LCMV Armstrong infection using a BD ARIA (BD). Total RNA was extracted from sorted cells using RNeasy Microkit (QIAGEN), and RNA was labeled with biotin with the BioArray High Yield RNA Transcript Labeling kit (Enzo Diagnostics) and purified with an RNeasy Mini kit (QIAGEN). The resulting cRNA was hybridized to GeneChip Mouse Gene 1.0 ST arrays at the UCSD VA/VMRF microarray and NGS core. The raw CEL files were obtained, and the data were normalized and analyzed with the GenePattern software suite.

ChIP and ChIP-seq. Methods for ChIP-Seq are described previously (Lin et al., 2010). Sequences were mapped to the mouse genome mm9 assembly (NCBI) using the Bowtie alignment tool and unique tags were visualized by preparing custom tracks for the UCSC Genome Browser where the total number of tags was normalized to 10⁷. DNA sequence analysis was performed using HOMER. Data are have been deposited in the the Gene Expression

Omnibus database under accession no. GSE46025. Foxo1 binding was referenced for T reg cells (Samstein et al., 2012) and two datasets for naive CD4 T cells (Naive 1; Ouyang et al., 2012; Naive2; this study). H3K4me3 histone marks were analyzed as previously described (Lin et al., 2010).

Statistical analysis. Differences between datasets were analyzed by unpaired two-tailed Student's *t* test or two-way ANOVA, as indicated.

This work was supported by National Institutes of Health grant 5R01AI103440 to SMH. The authors declare that they have no conflicting interests.

Submitted: 20 February 2013

Accepted: 6 May 2013

REFERENCES

- Almeida, M. 2011. Unraveling the role of FoxOs in bone—insights from mouse models. *Bone*. 49:319–327. <http://dx.doi.org/10.1016/j.bone.2011.05.023>
- Almeida, M., L. Han, M. Martin-Millan, C.A. O'Brien, and S.C. Manolagas. 2007. Oxidative stress antagonizes Wnt signaling in osteoblast precursors by diverting beta-catenin from T cell factor- to forkhead box O-mediated transcription. *J. Biol. Chem.* 282:27298–27305. <http://dx.doi.org/10.1074/jbc.M702811200>
- Araki, K., A.P. Turner, V.O. Shaffer, S. Gangappa, S.A. Keller, M.F. Bachmann, C.P. Larsen, and R. Ahmed. 2009. mTOR regulates memory CD8 T-cell differentiation. *Nature*. 460:108–112. <http://dx.doi.org/10.1038/nature08155>
- Bachmann, M.F., P. Wolint, S. Walton, K. Schwarz, and A. Oxenius. 2007. Differential role of IL-2R signaling for CD8+ T cell responses in acute and chronic viral infections. *Eur. J. Immunol.* 37:1502–1512. <http://dx.doi.org/10.1002/eji.200637023>
- Banerjee, A., S.M. Gordon, A.M. Intlekofer, M.A. Paley, E.C. Mooney, T. Lindsten, E.J. Wherry, and S.L. Reiner. 2010. Cutting edge: The transcription factor eomesodermin enables CD8+ T cells to compete for the memory cell niche. *J. Immunol.* 185:4988–4992. <http://dx.doi.org/10.4049/jimmunol.1002042>
- Barolo, S., and J.W. Posakony. 2002. Three habits of highly effective signaling pathways: principles of transcriptional control by developmental cell signaling. *Genes Dev.* 16:1167–1181. <http://dx.doi.org/10.1101/gad.976502>
- Best, J.A., D.A. Blair, J. Knell, E. Yang, V. Mayya, A. Doedens, M.L. Dustin, A.W. Goldrath; Immunological Genome Project Consortium. 2013. Transcriptional insights into the CD8(+) T cell response to infection and memory T cell formation. *Nat. Immunol.* 14:404–412. <http://dx.doi.org/10.1038/ni.2536>
- Beuneu, H., F. Lemaître, J. Deguine, H.D. Moreau, I. Bouvier, Z. Garcia, M.L. Albert, and P. Bousso. 2010. Visualizing the functional diversification of CD8+ T cell responses in lymph nodes. *Immunity*. 33:412–423. <http://dx.doi.org/10.1016/j.immuni.2010.08.016>
- Celli, S., Z. Garcia, H. Beuneu, and P. Bousso. 2008. Decoding the dynamics of T cell-dendritic cell interactions in vivo. *Immunol. Rev.* 221:182–187. <http://dx.doi.org/10.1111/j.1600-065X.2008.00588.x>
- Choo, D.K., K. Murali-Krishna, R. Anita, and R. Ahmed. 2010. Homeostatic turnover of virus-specific memory CD8 T cells occurs stochastically and is independent of CD4 T cell help. *J. Immunol.* 185:3436–3444. <http://dx.doi.org/10.4049/jimmunol.1001421>
- Curtsinger, J.M., C.M. Johnson, and M.F. Mescher. 2003. CD8 T cell clonal expansion and development of effector function require prolonged exposure to antigen, costimulation, and signal 3 cytokine. *J. Immunol.* 171:5165–5171.
- de la Torre, J.C., and M.B. Oldstone. 1992. Selective disruption of growth hormone transcription machinery by viral infection. *Proc. Natl. Acad. Sci. USA*. 89:9939–9943. <http://dx.doi.org/10.1073/pnas.89.20.9939>
- Fabre, S., F. Carrette, J. Chen, V. Lang, M. Semichon, C. Denoyelle, V. Lazar, N. Cagnard, A. Dubart-Kupferschmitt, M. Mangeney, et al. 2008. FOXO1 regulates L-Selectin and a network of human T cell homing molecules downstream of phosphatidylinositol 3-kinase. *J. Immunol.* 181:2980–2989.
- Gerlach, C., J.W. van Heijst, E. Swart, D. Sie, N. Armstrong, R.M. Kerkhoven, D. Zehn, M.J. Bevan, K. Schepers, and T.N. Schumacher. 2010. One naive T cell, multiple fates in CD8+ T cell differentiation. *J. Exp. Med.* 207:1235–1246. <http://dx.doi.org/10.1084/jem.20091175>
- Goodman, C.A., D.L. Mayhew, and T.A. Hornberger. 2011. Recent progress toward understanding the molecular mechanisms that regulate skeletal muscle mass. *Cell. Signal.* 23:1896–1906. <http://dx.doi.org/10.1016/j.cellsig.2011.07.013>
- Guo, K., J.E. McMinn, T. Ludwig, Y.H. Yu, G. Yang, L. Chen, D. Loh, C. Li, S.J. Chua Jr., and Y. Zhang. 2007. Disruption of peripheral leptin signaling in mice results in hyperleptinemia without associated metabolic abnormalities. *Endocrinology*. 148:3987–3997. <http://dx.doi.org/10.1210/en.2007-0261>
- Hand, T.W., W. Cui, Y.W. Jung, E. Sefik, N.S. Joshi, A. Chandele, Y. Liu, and S.M. Kaech. 2010. Differential effects of STAT5 and PI3K/AKT signaling on effector and memory CD8 T-cell survival. *Proc. Natl. Acad. Sci. USA*. 107:16601–16606. <http://dx.doi.org/10.1073/pnas.1003457107>
- Hedrick, S.M., R. Hess Michelini, A.L. Doedens, A.W. Goldrath, and E.L. Stone. 2012. FOXO transcription factors throughout T cell biology. *Nat. Rev. Immunol.* 12:649–661. <http://dx.doi.org/10.1038/nri3278>
- Hoogeboom, D., M.A. Essers, P.E. Polderman, E. Voets, L.M. Smits, and B.M. Burgering. 2008. Interaction of FOXO with beta-catenin inhibits beta-catenin/T cell factor activity. *J. Biol. Chem.* 283:9224–9230. <http://dx.doi.org/10.1074/jbc.M706638200>
- Hu, J.K., T. Kagari, J.M. Clingan, and M. Matloubian. 2011. Expression of chemokine receptor CXCR3 on T cells affects the balance between effector and memory CD8 T-cell generation. *Proc. Natl. Acad. Sci. USA*. 108:E118–E127. <http://dx.doi.org/10.1073/pnas.1101881108>
- Intlekofer, A.M., N. Takemoto, E.J. Wherry, S.A. Longworth, J.T. Northrup, V.R. Palanivel, A.C. Mullen, C.R. Gasink, S.M. Kaech, J.D. Miller, et al. 2005. Effector and memory CD8+ T cell fate coupled by T-bet and eomesodermin. *Nat. Immunol.* 6:1236–1244. <http://dx.doi.org/10.1038/ni1268>
- Jacob, J., and D. Baltimore. 1999. Modelling T-cell memory by genetic marking of memory T cells in vivo. *Nature*. 399:593–597. <http://dx.doi.org/10.1038/21208>
- Ji, Y., Z. Pos, M. Rao, C.A. Klebanoff, Z. Yu, M. Sukumar, R.N. Reger, D.C. Palmer, Z.A. Borman, P. Muranski, et al. 2011. Repression of the DNA-binding inhibitor Id3 by Blimp-1 limits the formation of memory CD8+ T cells. *Nat. Immunol.* 12:1230–1237. <http://dx.doi.org/10.1038/ni.2153>
- Joshi, N.S., W. Cui, A. Chandele, H.K. Lee, D.R. Urso, J. Hagan, L. Gapin, and S.M. Kaech. 2007. Inflammation directs memory precursor and short-lived effector CD8(+) T cell fates via the graded expression of T-bet transcription factor. *Immunity*. 27:281–295. <http://dx.doi.org/10.1016/j.immuni.2007.07.010>
- Kaech, S.M., and W. Cui. 2012. Transcriptional control of effector and memory CD8+ T cell differentiation. *Nat. Rev. Immunol.* 12:749–761. <http://dx.doi.org/10.1038/nri3307>
- Kaech, S.M., S. Hemby, E. Kersh, and R. Ahmed. 2002. Molecular and functional profiling of memory CD8 T cell differentiation. *Cell*. 111:837–851. [http://dx.doi.org/10.1016/S0092-8674\(02\)01139-X](http://dx.doi.org/10.1016/S0092-8674(02)01139-X)
- Kalia, V., S. Sarkar, S. Subramaniam, W.N. Haining, K.A. Smith, and R. Ahmed. 2010. Prolonged interleukin-2Ralpha expression on virus-specific CD8+ T cells favors terminal-effector differentiation in vivo. *Immunity*. 32:91–103. <http://dx.doi.org/10.1016/j.immuni.2009.11.010>
- Kallies, A., A. Xin, G.T. Belz, and S.L. Nutt. 2009. Blimp-1 transcription factor is required for the differentiation of effector CD8(+) T cells and memory responses. *Immunity*. 31:283–295. <http://dx.doi.org/10.1016/j.immuni.2009.06.021>
- Kerdiles, Y.M., D.R. Beisner, R. Tinoco, A.S. Dejean, D.H. Castrillon, R.A. DePinho, and S.M. Hedrick. 2009. Foxo1 links homing and survival of naive T cells by regulating L-selectin, CCR7 and interleukin 7 receptor. *Nat. Immunol.* 10:176–184. <http://dx.doi.org/10.1038/ni.1689>
- Kerdiles, Y.M., E.L. Stone, D.R. Beisner, M.A. McGargill, I.L. Ch'en, C. Stockmann, C.D. Katayama, and S.M. Hedrick. 2010. Foxo transcription factors control regulatory T cell development and function. *Immunity*. 33:890–904. <http://dx.doi.org/10.1016/j.immuni.2010.12.002>
- Kim, E.H., J.A. Sullivan, E.H. Plisch, M.M. Tejera, A. Jatzek, K.Y. Choi, and M. Suresh. 2012. Signal integration by Akt regulates CD8 T cell

- effector and memory differentiation. *J. Immunol.* 188:4305–4314. <http://dx.doi.org/10.4049/jimmunol.1103568>
- Kloet, D.E., and B.M. Burgering. 2011. The PKB/FOXO switch in aging and cancer. *Biochim. Biophys. Acta.* 1813:1926–1937. <http://dx.doi.org/10.1016/j.bbamcr.2011.04.003>
- Kousteni, S. 2012. FoxO1, the transcriptional chief of staff of energy metabolism. *Bone.* 50:437–443. <http://dx.doi.org/10.1016/j.bone.2011.06.034>
- Lin, Y.C., S. Jhunjhunwala, C. Benner, S. Heinz, E. Welinder, R. Mansson, M. Sigvardsson, J. Hagman, C.A. Espinoza, J. Dutkowski, et al. 2010. A global network of transcription factors, involving E2A, EBF1 and Foxo1, that orchestrates B cell fate. *Nat. Immunol.* 11:635–643. <http://dx.doi.org/10.1038/ni.1891>
- Melnik, B.C. 2011. Isotretinoin and FoxO1: A scientific hypothesis. *Dermatoendocrinol.* 3:141–165.
- Murali-Krishna, K., L.L. Lau, S. Sambhara, F. Lemonnier, J. Altman, and R. Ahmed. 1999. Persistence of memory CD8 T cells in MHC class I-deficient mice. *Science.* 286:1377–1381.
- Obar, J.J., M.J. Molloy, E.R. Jellison, T.A. Stoklasek, W. Zhang, E.J. Usherwood, and L. Lefrançois. 2010. CD4+ T cell regulation of CD25 expression controls development of short-lived effector CD8+ T cells in primary and secondary responses. *Proc. Natl. Acad. Sci. USA.* 107:193–198.
- Oldstone, M.B. 2002. Biology and pathogenesis of lymphocytic choriomeningitis virus infection. *Curr. Top. Microbiol. Immunol.* 263:83–117.
- Ouyang, W., W. Liao, C.T. Luo, N. Yin, M. Huse, M.V. Kim, M. Peng, P. Chan, Q. Ma, Y. Mo, et al. 2012. Novel Foxo1-dependent transcriptional programs control T(reg) cell function. *Nature.* 491:554–559.
- Paik, J.H., R. Kollipara, G. Chu, H. Ji, Y. Xiao, Z. Ding, L. Miao, Z. Tothova, J.W. Horner, D.R. Carrasco, et al. 2007. FoxOs are lineage-restricted redundant tumor suppressors and regulate endothelial cell homeostasis. *Cell.* 128:309–323.
- Parish, I.A., and S.M. Kaech. 2009. Diversity in CD8(+) T cell differentiation. *Curr. Opin. Immunol.* 21:291–297. <http://dx.doi.org/10.1016/j.coi.2009.05.008>
- Parretta, E., G. Cassese, A. Santoni, J. Guardiola, A. Vecchio, and F. Di Rosa. 2008. Kinetics of in vivo proliferation and death of memory and naive CD8 T cells: parameter estimation based on 5-bromo-2'-deoxyuridine incorporation in spleen, lymph nodes, and bone marrow. *J. Immunol.* 180:7230–7239.
- Pearce, E.L., and H. Shen. 2007. Generation of CD8 T cell memory is regulated by IL-12. *J. Immunol.* 179:2074–2081.
- Pearce, E.L., M.C. Walsh, P.J. Cejas, G.M. Harms, H. Shen, L.S. Wang, R.G. Jones, and Y. Choi. 2009. Enhancing CD8 T-cell memory by modulating fatty acid metabolism. *Nature.* 460:103–107. <http://dx.doi.org/10.1038/nature08097>
- Pekowska, A., T. Benoukraf, J. Zacarias-Cabeza, M. Belhocine, F. Koch, H. Holota, J. Imbert, J.C. Andrau, P. Ferrier, and S. Spicuglia. 2011. H3K4 tri-methylation provides an epigenetic signature of active enhancers. *EMBO J.* 30:4198–4210. <http://dx.doi.org/10.1038/emboj.2011.295>
- Pipkin, M.E., J.A. Sacks, F. Cruz-Guilloty, M.G. Lichtenheld, M.J. Bevan, and A. Rao. 2010. Interleukin-2 and inflammation induce distinct transcriptional programs that promote the differentiation of effector cytolytic T cells. *Immunity.* 32:79–90. <http://dx.doi.org/10.1016/j.immuni.2009.11.012>
- Rao, R.R., Q. Li, K. Odunsi, and P.A. Shrikant. 2010. The mTOR kinase determines effector versus memory CD8+ T cell fate by regulating the expression of transcription factors T-bet and Eomesodermin. *Immunity.* 32:67–78. <http://dx.doi.org/10.1016/j.immuni.2009.10.010>
- Rao, R.R., Q. Li, M.R. Gubbels Bupp, and P.A. Shrikant. 2012. Transcription factor Foxo1 represses T-bet-mediated effector functions and promotes memory CD8(+) T cell differentiation. *Immunity.* 36:374–387. <http://dx.doi.org/10.1016/j.immuni.2012.01.015>
- Rutishauser, R.L., and S.M. Kaech. 2010. Generating diversity: transcriptional regulation of effector and memory CD8 T-cell differentiation. *Immunol. Rev.* 235:219–233.
- Rutishauser, R.L., G.A. Martins, S. Kalachikov, A. Chande, I.A. Parish, E. Meffre, J. Jacob, K. Calame, and S.M. Kaech. 2009. Transcriptional repressor Blimp-1 promotes CD8(+) T cell terminal differentiation and represses the acquisition of central memory T cell properties. *Immunity.* 31:296–308. <http://dx.doi.org/10.1016/j.immuni.2009.05.014>
- Samstein, R.M., A. Arvey, S.Z. Josefowicz, X. Peng, A. Reynolds, R. Sandstrom, S. Neph, P. Sabo, J.M. Kim, W. Liao, et al. 2012. Foxp3 exploits a pre-existent enhancer landscape for regulatory T cell lineage specification. *Cell.* 151:153–166. <http://dx.doi.org/10.1016/j.cell.2012.06.053>
- Sarkar, S., V. Kalia, W.N. Haining, B.T. Konieczny, S. Subramaniam, and R. Ahmed. 2008. Functional and genomic profiling of effector CD8 T cell subsets with distinct memory fates. *J. Exp. Med.* 205:625–640. <http://dx.doi.org/10.1084/jem.20071641>
- Shin, H., S.D. Blackburn, A.M. Intlekofer, C. Kao, J.M. Angelosanto, S.L. Reiner, and E.J. Wherry. 2009. A role for the transcriptional repressor Blimp-1 in CD8(+) T cell exhaustion during chronic viral infection. *Immunity.* 31:309–320. <http://dx.doi.org/10.1016/j.immuni.2009.06.019>
- Stemberger, C., K.M. Huster, M. Koffler, F. Anderl, M. Schiemann, H. Wagner, and D.H. Busch. 2007. A single naive CD8+ T cell precursor can develop into diverse effector and memory subsets. *Immunity.* 27:985–997. <http://dx.doi.org/10.1016/j.immuni.2007.10.012>
- Sykes, S.M., S.W. Lane, L. Bullinger, D. Kalaitzidis, R. Yusuf, B. Saez, F. Ferraro, F. Mercier, H. Singh, K.M. Brumme, et al. 2011. AKT/FOXO signaling enforces reversible differentiation blockade in myeloid leukemias. *Cell.* 146:697–708. <http://dx.doi.org/10.1016/j.cell.2011.07.032>
- Takemoto, N., A.M. Intlekofer, J.T. Northrup, E.J. Wherry, and S.L. Reiner. 2006. Cutting Edge: IL-12 inversely regulates T-bet and eomesodermin expression during pathogen-induced CD8+ T cell differentiation. *J. Immunol.* 177:7515–7519.
- Tanchot, C., F.A. Lemonnier, B. Péramau, A.A. Freitas, and B. Rocha. 1997. Differential requirements for survival and proliferation of CD8 naive or memory T cells. *Science.* 276:2057–2062. <http://dx.doi.org/10.1126/science.276.5321.2057>
- Tothova, Z., R. Kollipara, B.J. Huntly, B.H. Lee, D.H. Castrillon, D.E. Cullen, E.P. McDowell, S. Lazo-Kallanian, I.R. Williams, C. Sears, et al. 2007. FoxOs are critical mediators of hematopoietic stem cell resistance to physiologic oxidative stress. *Cell.* 128:325–339. <http://dx.doi.org/10.1016/j.cell.2007.01.003>
- Williams, M.A., A.J. Tyznik, and M.J. Bevan. 2006. Interleukin-2 signals during priming are required for secondary expansion of CD8+ memory T cells. *Nature.* 441:890–893. <http://dx.doi.org/10.1038/nature04790>
- Wirth, T., and J.T. Harty. 2009. Initial TCR transgenic precursor frequency alters functional behaviour of CD8 T cells responding to acute infection. *Adv. Exp. Med. Biol.* 633:71–80. http://dx.doi.org/10.1007/978-0-387-79311-5_7
- Yang, C.Y., J.A. Best, J. Knell, E. Yang, A.D. Sheridan, A.K. Jesionek, H.S. Li, R.R. Rivera, K.C. Lind, L.M. D'Cruz, et al. 2011. The transcriptional regulators Id2 and Id3 control the formation of distinct memory CD8+ T cell subsets. *Nat. Immunol.* 12:1221–1229. <http://dx.doi.org/10.1038/ni.2158>
- Youngblood, B., E.J. Wherry, and R. Ahmed. 2012. Acquired transcriptional programming in functional and exhausted virus-specific CD8 T cells. *Curr. Opin. HIV AIDS.* 7:50–57. <http://dx.doi.org/10.1097/COH.0b013e32834ddcf2>
- Yu, Q., A. Sharma, and J.M. Sen. 2010. TCF1 and beta-catenin regulate T cell development and function. *Immunol. Res.* 47:45–55. <http://dx.doi.org/10.1007/s12026-009-8137-2>
- Zhang, N., and M.J. Bevan. 2011. CD8(+) T cells: foot soldiers of the immune system. *Immunity.* 35:161–168. <http://dx.doi.org/10.1016/j.immuni.2011.07.010>
- Zhang, X., S. Yalcin, D.F. Lee, T.Y. Yeh, S.M. Lee, J. Su, S.K. Mungamuri, P. Rimmelé, M. Kennedy, R. Sellers, et al. 2011. FOXO1 is an essential regulator of pluripotency in human embryonic stem cells. *Nat. Cell Biol.* 13:1092–1099. <http://dx.doi.org/10.1038/ncb2293>
- Zhou, X., and H.H. Xue. 2012. Cutting edge: generation of memory precursors and functional memory CD8+ T cells depends on T cell factor-1 and lymphoid enhancer-binding factor-1. *J. Immunol.* 189:2722–2726. <http://dx.doi.org/10.4049/jimmunol.1201150>
- Zhou, X., S. Yu, D.M. Zhao, J.T. Harty, V.P. Badovinac, and H.H. Xue. 2010. Differentiation and persistence of memory CD8(+) T cells depend on T cell factor 1. *Immunity.* 33:229–240. <http://dx.doi.org/10.1016/j.immuni.2010.08.002>